arXiv:1302.0803v1 [math.AG] 4 Feb 2013

THE MORI PROGRAM AND NON-FANO TORIC HOMOLOGICAL MIRROR SYMMETRY

MATTHEW BALLARD, COLIN DIEMER, DAVID FAVERO, LUDMIL KATZARKOV, AND GABRIEL KERR

Abstract. In the case of toric varieties, we continue the pursuit of Kontsevich's fundamental insight, Homological Mirror Symmetry, by unifying it with the Mori program. We give a refined conjectural version of Homological Mirror Symmetry relating semi-orthogonal decompositions of the B-model on toric varieties to semi-orthogonal decompositions on the A-model on the mirror Landau-Ginzburg models.

As evidence, we prove a new case of Homological Mirror Symmetry for a toric surface whose anticanonical bundle is not nef, namely a certain blow-up of $\mathbb{P}^2$ at three infinitesimally near points.

1. Introduction

In [AKO08], Auroux, Orlov, and the fourth named author discovered an "unusual" phenomenon regarding Homological Mirror Symmetry (HMS) for Hirzebruch surfaces. This phenomenon also arose, in greater generality, in [Abo09] where Abouzaid considered HMS for toric varieties whose anticanonical bundle is not ample.

The basic observation in the example of the Hirzebruch surface is the following. Consider the weighted projective stack,

$$\mathbb{P}(1:1:n) = [(\mathbb{A}^3 \setminus \{0\})/\mathbb{G}_m]$$

where $\mathbb{G}_m = \mathbb{C}^*$ acts with weights $1, 1, n$. The toric fan has three rays with primitive generators $(1,0)$, $(0,1)$, and $(-1,-n)$. Following the Givental-Hori-Vafa prescription [HV02], the mirror is a Landau-Ginzburg (LG) model

$$w : (\mathbb{C}^*)^2 \to \mathbb{C},$$

where

$$w(x,y) = x + y + \frac{1}{x^n y}.$$

Correspondingly, for the Hirzebruch surface

$$\mathbb{F}_n = \underline{\mathrm{Proj}}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(n)),$$

the (naive) mirror is a Landau-Ginzburg model

$$w' : (\mathbb{C}^*)^2 \to \mathbb{C},$$

where

$$w'(x,y) = x + y + \frac{1}{x} + \frac{1}{x^n y}.$$

It was observed in Lemma 5.3 of [AKO08] that for $n \geq 2$ these two functions are isotopic as exact Lefschetz fibrations and therefore have equivalent Fukaya-Seidel categories.

On the other hand, the bounded derived categories of coherent sheaves on $\mathbb{P}(1:1:n)$ and $\mathbb{F}_n$ are not equivalent for $n \neq 2$. For example, a full strong exceptional collection for $\mathrm{D}^{\mathrm{b}}(\mathrm{coh}\,\mathbb{F}_n)$ is given by

$$T = \mathcal{O}_{\mathbb{F}_n} \oplus \pi^* \mathcal{O}_{\mathbb{P}^1}(1) \oplus \mathcal{O}_{\pi}(1) \oplus \pi^* \mathcal{O}_{\mathbb{P}^1}(1) \otimes \mathcal{O}_{\pi}(1)$$

where $\pi : \mathbb{F}_n \to \mathbb{P}^1$ is the projection. Similarly, a full strong exceptional collection for $\mathrm{D}^{\mathrm{b}}(\mathrm{coh}\,\mathbb{P}(1:1:n))$ is given by

$$T' = \bigoplus_{i=0}^{n+2} \mathcal{O}_{\mathbb{P}(1:1:n)}(i).$$



FIGURE 1. Quiver representing the endomorphism algebra of T ($n = 4$)



FIGURE 2. Quiver representing the endomorphism algebra of T' ($n = 4$)

Figures 1 and 2 compare the endomorphism algebras of T and T' for $n = 4$ (where there are implicit relations $x_i x_j = x_j x_i$ when both sides of the equality have the same head and tail). Observe that removing the two central vertices from Figure 2 gives exactly Figure 1. In general, for $n \geq 2$ it is always the case that

$$\mathrm{End}(T) \cong \mathrm{End}(\mathcal{O}_{\mathbb{P}(1:1:n)} \oplus \mathcal{O}_{\mathbb{P}(1:1:n)}(1) \oplus \mathcal{O}_{\mathbb{P}(1:1:n)}(n) \oplus \mathcal{O}_{\mathbb{P}(1:1:n)}(n+1)).$$

This implies that for $n \geq 2$ there is a fully-faithful functor (see Theorem 2.29 of [AKO08])

$$\mathrm{MK}_n : \mathrm{D}^{\mathrm{b}}(\mathrm{coh}\,\mathbb{F}_n) \to \mathrm{D}^{\mathrm{b}}(\mathrm{coh}\,\mathbb{P}(1:1:n)). \tag{1.1}$$

In summary, while the Fukaya-Seidel categories of the LG mirrors are equivalent, the derived categories of coherent sheaves on the toric varieties are only related by a fully-faithful functor. It is natural to expect that the discrepancy comes from the fact that the anticanonical divisor on $\mathbb{F}_n$ is no longer nef for $n > 2$.

Moreover, in the mirror, this discrepancy can be accounted for through a certain deformation. Consider the family of isotopic LG models obtained by introducing the complex parameter b in the equation

$$w = x + y + \frac{1}{x} + \frac{b}{x^n y}.$$

For $b \neq 0$ these functions yield equivalent Fukaya-Seidel categories, and have $n+2$ critical points, in bijection with the $n+2$ indecomposible summands of T'. However, we have observed that $\mathrm{D^b}(\mathrm{coh}\,\mathbb{F}_n)$ corresponds to 4 of the indecomposible summands of T'. This is explained by Lemma 5.4 of [AKO08] which observes that if we let b go to zero, $n-2$ of these critical points go off towards infinity while 4 remain in a bounded region. Figure 3 is an image appearing in [AKO08]. It illustrates this phenomenon for $n = 8$. Therefore instead of the (naive) mirror, we can consider what we would call the homological mirror, which consists only of those critical fibers contained in the bounded region.

One aim of this work is to understand this phenomena conceptually in terms of moduli spaces of Landau-Ginzburg models, as introduced in [DKK12]. This work also introduced the notion of "radar screens" which gives a more general formalism for working with situations where critical loci of Landau-Ginzburg models assemble into regions upon a deformation.



FIGURE 3. The deformation $b \to 0$ $(n = 8)$

In general, mirror symmetry identifies complex moduli with Kähler moduli, and in the example above the complex deformation parameter b is mirror to a deformation of the complexified Kähler class. For a smooth toric variety X the Hodge numbers $h^{2,0}$ and $h^{0,2}$ vanish, hence symplectic forms can be identified with $\mathbb{R}$-divisors $\mathrm{H}^2(X, \mathbb{R}) \cong \mathrm{Pic}(X) \otimes \mathbb{R} =: \mathrm{Pic}_{\mathbb{R}}(X)$. Furthermore, the GKZ fan, which parametrizes the various GIT quotients obtained from the Cox quotient construction, can also be identified with a fan supported on the pseudo-effective cone in $\mathrm{Pic}_{\mathbb{R}}(X)$. In this way, degenerating the complex parameter of the mirror amounts to variation of GIT quotients (VGIT) on X. For the Hirzebruch surface, the two chambers of the GKZ fan correspond to $\mathbb{P}(1:1:n)$ and $\mathbb{F}_n$ as shown in Figure 4.



FIGURE 4. The GKZ fan of a Hirzebruch surface

In [BFK12], a precise relationship between the derived categories of quotient stacks coming from VGIT is given. It, in particular, reproduces the inclusion 1.1.

In this paper, we aim to entirely describe this phenomenon for a general run of the toric Mori program for toric projective DM stacks, building on themes from [DKK12] and [BFK12]. This includes a framework for Homological Mirror Symmetry for cases where the anticanonical divisor is not ample, nor even nef.

Indeed, in [DKK12], the authors demonstrated that any (straight line) run of the toric Mori program degenerates the mirror so that critical points can be arranged into successive annuli. This arrangement of critical points provides a semi-orthogonal decomposition of the Fukaya-Seidel category of this mirror.

Theorem 1 ([DKK12])**.** *For every maximal degenerated LG mirror ψ of a family $\psi(t)$ there is a semi-orthogonal decomposition*

$$\langle \overbrace{\mathrm{Fuk}^{\rightarrow}(\psi_1),\dots,\mathrm{Fuk}^{\rightarrow}(\psi_1)}^{-\mu_1},\dots,\overbrace{\mathrm{Fuk}^{\rightarrow}(\psi_s),\dots,\mathrm{Fuk}^{\rightarrow}(\psi_s)}^{-\mu_s}\rangle.$$

of the Fukaya-Seidel category $\mathrm{Fuk}^{\rightarrow}(\psi(t))$ *for* $t \approx 0$*. Here the ψ_i are regenerations of the degenerated LG model corresponding to each annulus and $-\mu_i$ are positive integers associated to the corresponding run of the toric Mori program.*

On the other side of mirror symmetry, we recall the following result.

Theorem 2 ([BFK12])**.** *Let X be a projective toric DM stack, equipped with a run of the toric Mori program. Starting from X, let $(W_1,\mu_1),\dots,(W_s,\mu_s)$ be the wall data associated to the run. Then there exists a semi-orthogonal decomposition*

$$\langle \overbrace{\mathrm{D^b}(\mathrm{coh}\, W_1),\dots,\mathrm{D^b}(\mathrm{coh}\, W_1)}^{-\mu_1},\dots,\overbrace{\mathrm{D^b}(\mathrm{coh}\, W_s),\dots,\mathrm{D^b}(\mathrm{coh}\, W_s)}^{-\mu_s}\rangle.$$

of $\mathrm{D^b}(\mathrm{coh}\, X)$.

In conclusion, a (straight line) run of the toric Mori program for X yields both a semi-orthogonal decomposition of $\mathrm{D^b}(\mathrm{coh}\, X)$ and a semi-orthogonal decomposition of the Fukaya-Seidel category of the Landau-Ginzburg mirror of X.

From here on, let us say that a variety is **nef-Fano** if it has $\mathbb{Q}$-factorial singularities and the anticanonical divisor $-K_X$ is big and nef. This condition is also often called weak Fano, usually with the additional assumption that the singularities are canonical. We will say that a Deligne-Mumford (or DM) stack is nef-Fano if its underlying coarse variety is nef-Fano.

Given X there is always a (straight line) run of the toric Mori program which contains a nef-Fano projective toric DM stack X' as a phase (see Proposition 2.3.11). This allows us

to view the mirror to any X as a Landau-Ginzburg model coming from a neighborhood of some collection of critical points in the mirror to X'. Therefore we propose that, like the homological mirror to $\mathbb{F}_n$, the homological mirror of any projective toric DM stack X can be realized as a regeneration of the degenerated Landau-Ginzburg mirror of X' (in the sense of [DKK12]).

With this picture in mind, we extend one half of Homological Mirror Symmetry for toric varieties as follows:

Conjecture 1. *Given, γ, a straight line run of the toric Mori program for a projective DM toric stack, X, and its associated maximal degeneration ψ of an LG mirror pencil, let*

$$\mathrm{Fuk}^{\rightarrow}(\psi(t)) = \langle \mathcal{T}_1, \ldots, \mathcal{T}_r \rangle$$
$$\mathrm{D^b}(\mathrm{coh}\, X) = \langle \mathcal{S}_1, \ldots, \mathcal{S}_r \rangle$$

be the semi-orthogonal decompositions associated to ψ and γ.

Then there exists an equivalence of triangulated categories

$$\Phi_\psi : \mathrm{Fuk}^{\rightarrow}(\psi(t)) \to \mathrm{D^b}(\mathrm{coh}\, X)$$

which restricts to equivalences $\Psi_\psi : \mathcal{T}_i \to \mathcal{S}_i$ for all $1 \leq i \leq r$.

As evidence, we prove Conjecture 1 in the following case: take $\mathbb{P}^2$, blow it up at a point, and then blow up at the two points on the exceptional locus corresponding to the two coordinate lines passing through the original point. Call the resulting space X.

Theorem 3. *Conjecture 1 is true for a choice of run of the Mori program on X. In particular, Homological Mirror Symmetry holds for X.*

1.1. **Relation with other work.** Homological Mirror Symmetry (HMS) was conceived for Calabi-Yau varieties and for toric varieties by Kontsevich [Kon95, Kon98].

Seidel proved HMS for the projective plane [Sei00] and Auroux, Orlov, and the fourth named author proved HMS for weighted projective planes [AKO08] and del Pezzo surfaces (toric and non-toric) [AKO06]. Independently, Ueda proved HMS for toric del Pezzo surfaces [Ued06] and Ueda together with Yamazaki then proved it for toric orbifolds of del Pezzo surfaces [UY07]. Furthermore, Bondal and Ruan have announced a proof for weighted projective spaces of all dimensions [BR].

Abouzaid proved HMS for smooth Fano toric varieties in [Abo09] and, for general smooth projective toric varieties, provided an embedding

$$\mathrm{D^b}(\mathrm{coh}\, X) \to \mathrm{Fuk}^{\rightarrow}(w)$$

where w is the Hori-Vafa Landau-Ginzburg mirror. We emphasize that when X is not Fano, the image of Abouzaid's functor is not well understood in terms of mirror symmetry. Fang, Liu, Treumann, and Zaslow proved a very similar statement via different means in [FLTZ08]. A natural question to ask, given the results of [Abo09] and [FLTZ08], is: for a non-nef Fano projective toric DM stack X, how does one naturally determine the subcategory of the Fukaya-Seidal category of the Hori-Vafa mirror which corresponds to the derived category of X? This paper proposes that following the toric Mori program through mirror symmetry gives the answer.

In the case of Picard rank one, Conjecture 1 reduces to the usual statement of Homological Mirror Symmetry identically. For higher Picard rank, it includes the usual HMS statement, with additional structure.

While we are concerned with comparing the B-model on the toric stack to the A-model on the mirror LG model, there has also been work in the other direction, comparing the A-model on the toric stack to the B-model on the LG model. For $\mathbb{P}^2$, Chan and Leung showed how to produce matrix factorizations via Strominger-Yau-Zaslow [SYZ96] fibrations. In forthcoming work [AFOOO], Abouzaid, Fukaya, Oh, Ohta, and Ono advance further this direction of HMS. Abouzaid and Auroux have pointed out that our mirrors do not suffice for this other direction of mirror symmetry.

A similar result to Theorem 2 can be proven using Kawamata's pioneering work [Kaw06]. The connection between the Mori program and derived categories also goes back to Kawamata in [Kaw05, Kaw02] and extended in [BFK12] and [H-L12].

On the other hand, as we will see, the moduli space of LG models (and it's compactification) is a crucial ingredient for the A-side of our approach. Work in preparation of Kontsevich, Pantev, and the fourth named author suggests a definition of a moduli space of LG models for essentially arbitrary potentials [KKP13]. In another direction, it was pointed out to us by Keel that the results of [GKH11] produce mirror families for a large class of rational surfaces which carry over many of the critical features we make use of for the case of toric mirrors.

In future work we intend to study the program suggested in this work for these rational surfaces. In summary, we anticipate that the conjectural approach to proving HMS we outline here will find applicability for more general classes of varieties.

1.2. **Outline.** Let us briefly summarize the structure of this paper. We begin in Section 2.3 with a discussion of necessary background, including categorical and homological formalities and the general setup of the toric Mori program. In Section 3, we discuss the behavior of derived categories under variation of GIT and the results of [BFK12], with an emphasis on the connections with Mori theory. Section 4 discusses the other side of mirror symmetry with an overview of [DKK12] and general machinery for degenerations of mirror toric LG models. We conclude in Section 5 with a conjectural program towards a proof of Conjecture 1, and a complete execution of the program for a new example.

Acknowledgments: The authors have benefited immensely from conversations and correspondence with Denis Auroux, R. Paul Horja, Dmitri Orlov, Alexei Bondal, Kentaro Hori, Dragos Deliu, Umut Isik, Mohammed Abouzaid, Sean Keel, Dmitri Orlov, Alexander Kuznetsov, Yan Soibelman, and Maxim Kontsevich and would like to thank them all for their time, patience, and insight. The authors were funded by NSF DMS 0854977 FRG, NSF DMS 0600800, NSF DMS 0652633 FRG, NSF DMS 0854977, NSF DMS 0901330, FWF P 24572 N25, by FWF P20778 and by an ERC Grant. The first author was funded, in addition, by NSF DMS 0838210 RTG.

2. Background

2.1. **A_∞-algebras and homological perturbation.** We will assume that the reader is acquainted with the basics of A_∞-algebras, including modules, morphisms, and derived categories. Excellent references for the material are [Sta63, Kel01, L-H03]. We begin by recalling the most important nontrivial equivalence relation on A_∞-algebras.

Definition 2.1.1. A morphism of A_∞-algebras, $f : (A, m_n^A) \to (B, m_n^B)$, is called a **quasi-isomorphism** if the morphism on cohomology,

$$\mathrm{H}^*(f) : \mathrm{H}^*(A, m_1^A) \to \mathrm{H}^*(B, m_1^B),$$

is an isomorphism.

Quasi-isomorphic A_∞-algebras have equivalent derived categories of A_∞-modules. For a proof, see [L-H03].

Lemma 2.1.2. *If $f : (A, m_n^A) \to (B, m_n^B)$ is quasi-isomorphism, then there is an equivalence,*

$$\mathrm{D}(\mathrm{perf}\, A) \cong \mathrm{D}(\mathrm{perf}\, B),$$

between the derived categories of perfect modules.

Isomorphism classes of A_∞-algebras can be quite large. Quasi-isomorphism classes will be even larger. We will be interested in locating a managable representative of a quasi-isomorphism equivalence class as we will need to perform full computations of the A_∞-structure.

One particularly useful class of A_∞-algebras is the following.

Definition 2.1.3. An A_∞-algebra, (A, m_n^A), is called **minimal** if $m_1 = 0$.

Given a dg-algebra $(\widehat{A}, d, m)$, let A denote its cohomology algebra. Suppose we have a collection of morphisms of vector spaces,

$$h \circlearrowright \widehat{A} \underset{i}{\overset{\pi}{\rightleftarrows}} A$$

satisfying

$$\pi \circ i = 1 \text{ and } i \circ \pi = 1 - dh - hd \text{ and } h^2 = 0.$$

Homological perturbation, see e.g. [Mar04], gives explicit higher products m_n^A on A and a quasi-isomorphism between $\widehat{A}$ and (A, m_n^A). The formula for m_n^A is given as sum over trees with n leaves such that the valency of any internal vertex is 3. This is described now.

Given a tree, T, oriented from leaves to root with n leaves, we write, $m_{T,n}^A$, for the composition of operators induced by the following rules:

- For each leaf compose with the map $i : A \to \widehat{A}$
- Apply multiplication at every internal vertex
- For each internal edge compose with the map $h : \widehat{A} \to \widehat{A}$
- Apply $\pi : \widehat{A} \to A$ at the root

The m_n^A are given by the formula

$$m_n^A = \sum_T (-1)^{\vartheta(T)} m_{T,n}^A \tag{2.1}$$

where the sum is taken over all planar trees with n-leaves. For the precise signs, $\vartheta(T)$, see [Mar04]. We will only need the formula for $n = 3$ which we display graphically.



In terms of equations, we have

$$m_3^A(a,b,c) = \pi(m(h(m(i(a),i(b))),i(c))) - (-1)^{|a|}\pi(m(i(a),h(m(i(b),i(c))))) \tag{2.2}$$

where $|a|$ is the degree of a.

2.2. **Exceptional collections, quivers, and equivalences.** Given a triangulated category, we often wish to describe it by decomposing it into more manageable subcategories. To this end, a common structure appearing in [BFK12] and [DKK12] is that of a semi-orthogonal decomposition. In this section we quickly review this notion and focus on the special case of a full exceptional collection. Standard references for the material are [Bon89, BK89, Kel01].

Definition 2.2.1. A **semi-orthogonal decomposition** of a triangulated category, $\mathcal{T}$, is a sequence of full triangulated subcategories, $\mathcal{A}_1, \ldots, \mathcal{A}_m$, in $\mathcal{T}$ such that $\mathcal{A}_i \subset \mathcal{A}_j^{\perp}$ for $i < j$ and, for every object $T \in \mathcal{T}$, there exists a diagram:



where all triangles are distinguished and $A_k \in \mathcal{A}_k$. We shall denote a semi-orthogonal decomposition by $\langle \mathcal{A}_1, \ldots, \mathcal{A}_m \rangle$. If the subcategories in a semi-orthogonal decomposition are the essential images of fully-faithful functors, $\Upsilon_i : \mathcal{A}_i \to \mathcal{T}$, we shall also denote a semi-orthogonal decomposition by

$$\langle \Upsilon_1, \ldots, \Upsilon_m \rangle.$$

Assume that $\mathcal{T}$ is linear over a field, k. Let $E_1, \ldots, E_n$ be objects of $\mathcal{T}$. We say that $E_1, \ldots, E_n$ is an **exceptional collection** if

$$\mathrm{Hom}_{\mathcal{T}}(E_i, E_i[l]) = \begin{cases} k & \text{if } l = 0 \\ 0 & \text{otherwise} \end{cases}$$

for all i and

$$\mathrm{Hom}_{\mathcal{T}}(E_j, E_i[l]) = 0, \forall j > i, \forall l.$$

We say that $E_1, \ldots, E_n$ is **full** if $E_1, \ldots, E_n$ generates $\mathcal{T}$. We say $E_1, \ldots, E_n$ is a **strong** exceptional collection if

$$\mathrm{Hom}_{\mathcal{T}}(E_i, E_j[l]) = 0, l \neq 0, \forall i, j.$$

Possession of a full exceptional collection allows for a simplified presentation of an (enhanced) triangulated category. Let $E_1, \ldots, E_n$ be a full exceptional collection in $\mathcal{T}$. Then to $E_1, \ldots, E_n$ we can associate a graded quiver with relations, (Q, R). The vertex set of Q is the set $\{1, 2, \ldots, n\}$. Take a minimal set, S, of generators for the graded endomorphism algebra

$$A = \bigoplus_{l \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{T}}(\oplus_{j=1}^{n} E_j, \oplus_{j=1}^{n} E_j[l])$$

consisting of elements of the morphism spaces $\mathrm{Hom}_{\mathcal{T}}(E_i, E_j[l])$. The arrows from i to j in Q will be the elements of S lying in $\mathrm{Hom}_{\mathcal{T}}(E_i, E_j[l])$ for some $l \in \mathbb{Z}$. The degree of the arrow will be l.

To determine R, we note that, as we used a set of generators for A, there is a natural surjection

$$\pi : \mathbb{C}\, Q \to A$$

from the path algebra of Q to A. The ideal of relations, R, is the kernel of π.

One would hope that $\mathcal{T}$ is equivalent to the derived category of (Q, R) or equivalently to the derived category of graded A-modules. However, this hope is too naive. One must assume that $\mathcal{T}$ possesses some additional structure, an enhancement [BK89], to get something close to this.

Proposition 2.2.2. *If $\mathcal{T}$ possesses an enhancement, then A can be given the structure of a minimal A_∞-algebra, also denoted by A, such that there is an equivalence*

$$\mathcal{T} \cong \mathrm{D}(\mathrm{perf}\, A)$$

between $\mathcal{T}$ and the derived category of perfect A_∞-modules over the A_∞-algebra, A.

For a proof of the proposition see [Bon89, Kel94].

The above proposition allows us to determine $\mathcal{T}$ up to equivalence from a finite set of data: the graded quiver Q, the relations, R, and the higher A_∞ maps,

$$m_n^A : A^{\otimes n} \to A, \ n \geq 3.$$

Let us recall a well-known example: the Beilinson exceptional collection [Bei78]

$$\mathcal{O}_{\mathbb{P}^n}, \mathcal{O}_{\mathbb{P}^n}(1), \ldots, \mathcal{O}_{\mathbb{P}^n}(n)$$

is a strong full exceptional collection for the derived category $\mathrm{D}^{\mathrm{b}}(\mathrm{coh}\, \mathbb{P}^n)$. The graded quiver we get is pictured in Figure 5. All arrows are in degree 0.



FIGURE 5. Beilinson quiver

The ideal of relations is generated by

$$a_{i+1j}a_{ik} - a_{i+1k}a_{ij}$$

for $0 \leq i \leq n-1$ and $0 \leq j,k \leq n$. It expresses commutativity of the polynomial ring. Finally, the higher A_∞-structure maps are trivial for degree reasons.

Thus, whenever we can find an enhanced triangulated category, $\mathcal{T}$, and $(n+1)$-objects, $T_0, \ldots, T_n$ of $\mathcal{T}$ such that the endomorphism algebra of $\oplus_{i=0}^n T_i$ is the path algebra of the Beilinson quiver modulo the commutativity relations, we automatically get a fully-faithful functor,

$$\mathrm{D}^{\mathrm{b}}(\mathrm{coh}\,\mathbb{P}^n) \to \mathcal{T}.$$

If we can show that $\mathcal{T}$ generates, we will get an equivalence. Note that this phenomenon explains why the quivers from Figures 1 and 2 yield the fully-faithful functor MK_n in (1.1).

2.3. **The toric Mori program.** In this section, we briefly review the toric Mori program, also called the toric minimal model program, which originated in the work of Reid [Rei83]; standard references are the books of Cox, Little, Schenck [CLS11, Chapters 14-15] and Matsuki [Mat02, Chapter 14]. The subject has interpretations in terms of birational geometry, variation of Geometric Invariant Theory (VGIT), and combinatorial modifications of toric fans. It can be developed in each language in isolation.

We will focus mainly on VGIT. We choose this direction for its immediate relation with the work of [BFK12] and for its simplicity of presentation. For the birational geometer interested in comparing our presentation through VGIT to their more familiar perspective of flips, flops, and Mori contractions, we recommend reviewing [HK00]. For those more comfortable with the combinatorial viewpoint, we recommend [CLS11]. However, to make connections with [DKK12], we will also need to recall some combinatorial properties. We begin by recalling the general setup of VGIT [DH98, Tha96].

Let G be a linearly reductive group acting on a variety X which is equipped with a G-equivariant line bundle $\mathcal{L}$. Following Mumford [MFK94], we consider three subsets of X:

Definition 2.3.1.

$$\begin{aligned} X^{\mathrm{ss}}(\mathcal{L}) &= \{x \in X \mid \exists f \in \mathrm{H}^0(X, \mathcal{L}^n)^G \text{ with } n > 0, f(x) \neq 0, \text{ and } X_f \text{ affine}\} \\ X^{\mathrm{s}}(\mathcal{L}) &= \{x \in X^{\mathrm{ss}}(\mathcal{L}) \mid G \cdot x \text{ is closed in } X^{\mathrm{ss}}(\mathcal{L}) \text{ and } G_x \text{ is finite}\} \\ X^{\mathrm{us}}(\mathcal{L}) &= X \setminus X^{\mathrm{ss}}(\mathcal{L}). \end{aligned}$$

where $X_f = \{x \in X \mid f(x) \neq 0\}$. The subsets $X^{\mathrm{ss}}(\mathcal{L}), X^{\mathrm{s}}(\mathcal{L})$, and $X^{\mathrm{us}}(\mathcal{L})$, are called the **semi-stable, stable**, and **unstable locus**, respectively, of X. Each subset is naturally a subvariety of X.

Definition 2.3.2. With notation as above, the **GIT quotient** of X by G with respect to $\mathcal{L}$ is the quotient stack, $[X^{\mathrm{ss}}(\mathcal{L})/G]$. We denote the GIT quotient by $X/\!\!/_{\mathcal{L}}G$.

The reader should note that this is *not* the definition of a GIT quotient given in [MFK94], as we work with stacks. Indeed, the GIT quotient appearing in [MFK94] is the coarse moduli space of the GIT quotient appearing here, assuming that the stack is Deligne-Mumford.

The GIT quotient depends on the choice of $\mathcal{L}$. Exploring the relationship between the quotient stacks resulting from different choices of $\mathcal{L}$ has deep ties with birational geometry. Let $\mathrm{Pic}^G(X)$ be the group of G-equivariant line bundles on X. The associated vector space, $\mathrm{Pic}^G_{\mathbb{R}}(X)$, naturally forms a parameter space for GIT quotients. We say that $\mathcal{L}$ is G-**effective** if $\mathcal{L}$ is ample and $X^{\mathrm{ss}}(\mathcal{L}) \neq \emptyset$ and we let $\mathrm{Eff}^G_{\mathbb{R}}(X)$ denote the cone of G-effective equivariant line bundles.

Definition 2.3.3. The **GIT fan** for the action of G on X is a fan in $\mathrm{Pic}^G_{\mathbb{R}}(X)$ whose support is $\mathrm{Eff}^G_{\mathbb{R}}(X)$. The GIT fan is the unique fan characterized by the following property: the interiors of the cones of the GIT fan are exactly the subsets of constant semi-stable locus.

In general, the GIT fan may not exist. However, in many cases, it does. For example, if X is proper [DH98, Tha96, Res00] it exists. This implies that it exists for linear G-actions on $X = \mathbb{A}^n$ since we can compactify $\mathbb{A}^n$ to $\mathbb{P}^n$ and extend the action correspondingly.

We now leave the general setting of GIT to focus on toric setting: $X = \mathbb{A}^n$ and G is a subset of $\mathbb{G}_m^n$. Affine space, $\mathbb{A}^n$, has trivial Picard group so the set of G-equivariant line bundles is in bijection with characters of the group G, denoted by $\widehat{G}$. Given a character, χ, of G, we denote the corresponding G-equivariant line bundle by $\mathcal{O}(\chi)$ and the corresponding GIT quotient by $\mathbb{A}^n /\!\!/_\chi G$.

Definition 2.3.4. When $G \subseteq \mathbb{G}_m^n$ is a subgroup of an algebraic torus, we shall call a GIT quotient, $\mathbb{A}^n /\!\!/_\chi G$, of affine space $\mathbb{A}^n$ by G, a **toric stack**. We will say that the quotient is **projective** if $\mathbb{C}[x_1, \ldots, x_n]^G = \mathbb{C}$. If the quotient is represented by a scheme, we shall call the quotient a **toric variety**.

Remark 2.3.5. We have already mentioned that the GIT fan for toric actions on affine space exists. It is straightforward to see that a toric stack is Deligne-Mumford if and only if χ lies in the interior of a top dimensional cone of the GIT fan. This is equivalent to the underlying coarse toric variety being simplicial.

Remark 2.3.6. Note that, in general, a GIT quotient of affine space is automatically projective over an affine variety. In particular, if the quotient is proper, then it is projective; this occurs when $\mathbb{C}[x_1, \ldots, x_n]^G = \mathbb{C}$. However, the most common definition of a toric stack proceeds through a stacky fan, as in [BCS05] or the more general construction in [GS11]. These approaches also include non-projective toric stacks. At least in the case of varieties, our more restrictive definition has appeared elsewhere in the literature, see [MiSt05] as an example.

GIT fans for toric varieties have deeper roots. For toric varieties, when $\mathbb{A}^n$ is the spectrum of the Cox ring, the GIT fan was first considered by Oda-Park [OP91], motivated by a dual construction for polytopes introduced by Gel'fand, Kapranov, and Zelevinsky [GKZ94]. As such, it often goes by other names, including the **GKZ fan** or **secondary fan**, and it admits a purely combinatorial description, again see [CLS11, Mat02] for details.

We briefly recall the structure of the GKZ fan for toric quotients, again see [DH98, Tha96] for a treatment of the general case. The inclusion, $G \subset \mathbb{G}_m^n$ yields an exact sequence of abelian groups

$$0 \to M \xrightarrow{\delta} \mathbb{Z}^n \xrightarrow{\gamma} \widehat{G} \to 0, \tag{2.3}$$

of abelian groups, where M denotes the kernel of γ. Tensoring with $\mathbb{R}$, we obtain a sequence of vector spaces

$$0 \to M_{\mathbb{R}} \to \mathbb{R}^n \xrightarrow{\gamma_{\mathbb{R}}} \widehat{G}_{\mathbb{R}} \to 0.$$

Let $\beta_i := \gamma(e_i)$ with $e_1, \ldots, e_n$ the standard basis of $\mathbb{Z}^n$. Let C_β be the cone in $\widehat{G}_{\mathbb{R}}$ spanned by $\beta_1, \ldots, \beta_n$. It is easy to check that the characters in C_β are exactly those whose corresponding GIT quotients are non-empty.

Remark 2.3.7. In addition to the combinatorial descriptions, this fan also admits descriptions in terms of birational geometry. Recall that we can identify $\widehat{G}_{\mathbb{R}}$ with $\mathrm{Pic}_{\mathbb{R}}(X)$ when

X is a projective simplicial toric variety representing the corresponding quotient. In turn, $\mathrm{Pic}_{\mathbb{R}}(X)$ coincides in this case with $N^1(X)$, the space of divisors modulo numerical equivalence. The cone C_β then coincides with the cone of effective divisors $\mathrm{Eff}(X)$. This cone admits a fan called the Mori fan, where divisors in the relative interior of the same chamber define the same rational contraction; see e.g. [HK00] for a thorough discussion which in particular proves the equivalence of this fan with the GIT fan. We will occasionally use the standard facts that the cone $\mathrm{Amp}(X) \subset N^1(X)$ of ample divisors is the relative interior of the closed cone of nef divisors $\mathrm{Nef}(X)$, that the relative interior of the cone $\mathrm{Eff}(X)$ is the cone of big divisors $\mathrm{Big}(X)$, and that in the toric case $\mathrm{Eff}(X)$ is closed.

The maximal cones in the GIT fan are called **chambers**, and the codimension 1 cones are called **walls**. We say that two smooth toric projective DM stacks are **neighbors**, if there is a GKZ fan $\mathcal{F}_{\mathrm{GKZ}}$ such that the two stacks are GIT quotients corresponding to adjacent chambers separated by a wall. Motivated by string theory terminology, given a fixed GKZ fan, if X occurs as a GIT quotient corresponding to some maximal chamber, we say X is a **phase** of $\mathcal{F}_{\mathrm{GKZ}}$.

Now, let C_+ and C_- be two adjacent chambers in a GKZ fan. Let τ be the wall defined by the intersection of C_+ and C_-. The wall τ spans a linear hyperplane, H. Let $\lambda : \mathbb{G}_m \to G$ be a primitive one-parameter subgroup whose kernel, once tensored with $\mathbb{R}$, defines H. Furthermore, choose λ so that the pairing of λ with any character in C_+ is nonnegative. Set

$$G_\lambda := G/\lambda(\mathbb{G}_m).$$

Let α be a character in the interior of τ. Finally, consider the GIT quotient

$$W := (\mathbb{A}^n)^\lambda /\!\!/_\alpha G_\lambda$$

of the fixed locus of λ by G_λ via $\mathcal{O}(\alpha)$.

Set

$$\mu := -\langle \lambda, K_{[\mathbb{A}^n/G]} \rangle$$

where $K_{[\mathbb{A}^n/G]} = -\sum_i \beta_i$.

Definition 2.3.8. For a wall τ separating adjacent chambers C_+ and C_- in a GIT fan, we call (W, μ) the **wall data** associated to τ.

We are interested not in only in individual walls, but sequences of walls and the corresponding sequences of maximal chambers. Of particular importance are sequences which comes from a sequence of Mori operations (i.e. flops, directed flips, divisorial contractions, and Mori fibrations). To this end, fix a projective toric DM stack X and consider a continuous path $\gamma : [0, 1] \to \widehat{G}_{\mathbb{R}}$ such that:

- $\gamma(0)$ lies in the ample cone of X,
- $\gamma(1)$ lies outside the support of the GKZ fan,
- γ does not pass through any cones in the GKZ fan of codimension 2.

For any $0 < t < 1$, if $\gamma(t)$ lies in a wall, then we can consider the corresponding adjacent chambers C_+ and C_- where $\gamma(t - \epsilon) \in C_-$ and $\gamma(t + \epsilon) \in C_+$ for sufficiently small ϵ. This allows us to define a sequence of wall data along the path.

Definition 2.3.9. A path, γ, as above, is called a **run of the toric Mori program** if whenever γ intersects a wall τ one has $\mu \leq 0$ for the wall data (W, μ). If γ is run of the toric Mori program and is a line segment with respect to the Euclidean topology on $\widehat{G}_{\mathbb{R}}$ we call γ a **straight line run**.

We will frequently make use of the following elementary observation, which essentially states that any phase can be reached by a straight line run.

Lemma 2.3.10. *Let X be a projective toric DM stack with GKZ fan $\mathcal{F}_{\mathrm{GKZ}}$. Let Y be any phase of $\mathcal{F}_{\mathrm{GKZ}}$ with corresponding chamber C_Y. Fix any character χ in the ample cone of X. There is a line segment γ with respect to the Euclidean topology on $\widehat{G}_{\mathbb{R}}$ satisfying the three conditions above such that $\gamma(0) = \chi$ and $\gamma(a)$ lies in C_Y for some $0 < a < 1$.*

Proof. Fix the Euclidean metric on $\widehat{G}_{\mathbb{R}}$ and take $\epsilon > 0$ sufficiently small so that the sphere $S_\epsilon(\chi)$ of radius ϵ centered at χ is contained in the ample cone. Any point $\alpha \in S_\epsilon(\chi)$ determines a unique ray emanating from χ. Consider the collection of such points which determine a ray which eventually passes outside the pseudo-effective cone. As X is projective, the psedo-effective cone is strongly convex. Hence, the collection of all such points with a ray hitting the interior of C_Y is a non-empty open subset of the the sphere as C_Y is of maximum dimension. However, the locus of all points whose ray intersects a codimension 2 cone of $\mathcal{F}_{\mathrm{GKZ}}$ is a cell complex of real codimension at least 1. □

As discussed in the introduction, one of the aims of this paper is to study Homological Mirror Symmetry for smooth toric DM stacks which are not Fano (or nef-Fano). This following simple observation lets us always reach this case by a straight line run.

Proposition 2.3.11. *Let X be a projective toric DM stack. Then there exists a nef-Fano projective toric DM stack X' and a straight line run γ in the GKZ fan of X which starts at the phase X' and passes through the phase X and its way to the exterior of the pseudo-effective cone. Moreover, if the anticanonical divisor $-K_X$ is contained in the relative interior of a maximal chamber of the GKZ fan of X, then X' may be taken to be Fano.*

Proof. Let C_{-K} be any chamber which contains $-K_X$ (possibly in its boundary), and let X' be GIT quotient corresponding to this chamber. By the lemma above, let γ be a line segment starting in the relative interior of C_{-K}, passing through the ample cone of X, and eventually exiting the pseudo-effective cone.

To see that this path is a run of the toric Mori program, consider a value $0 < a < 1$ such that $\gamma(a)$ lies in a wall of the GKZ fan of X. By definition,

$$\mu = -\langle \lambda, K_{[\mathbb{A}^n/G]} \rangle$$

where λ is a one-parameter subgroup with $\langle \lambda, \gamma(a+\epsilon) \rangle > 0$ and $\langle \lambda, \gamma(a-\epsilon) \rangle < 0$ for some sufficiently small value of $\epsilon > 0$. Now, since our path is a straight line we have $\langle \lambda, \gamma(a-\epsilon) \rangle < 0$ for any $\epsilon > 0$. In particular, $\langle \lambda, \gamma(0) \rangle < 0$ and remains negative on the relative interior of C_{-K}. By continuity $\langle \lambda, v \rangle \leq 0$ for any $v \in C_{-K}$. Hence $\mu = -\langle \lambda, K_{[\mathbb{A}^n/G]} \rangle \leq 0$ as desired.

Recall that the anticanonical divisor on a projective toric variety is automatically effective and big. To see that it is also nef, notice that the GKZ fan of X' is a projection of the GKZ fan of X (possibly the identity projection), and the nef cone of X' is the image of C_{-K} under this projection, from which the proposition immediately follows. □

When we consider toric mirror symmetry, it will be natural to consider an enlargement of the GKZ fan which is complete. The following proposition is well-known.

Proposition 2.3.12. *Let X be a projective toric DM stack. Then the GKZ fan of X is naturally a subfan of the GKZ fan of the total space of the anticanonical bundle* $\mathrm{Tot}(-K_X)$*, and moreover the GKZ fan of* $\mathrm{Tot}(-K_X)$ *is complete.*

Following [AGM94], we call the GKZ fan of the anticanonical bundle of X the **enlarged GKZ fan** of X. That this enlarged fan is relevant for mirror symmetry is reasonable, as $\mathrm{Tot}(-K_X)$ is log Calabi-Yau.

3. The B-side

3.1. **The toric Mori program and derived categories.** The purpose of the Mori program is to take a projective variety, X, and, by a controlled sequence of birational morphisms, produce a "simplest" birational model of X. As birational geometry and derived categories are intimately tied, it is natural to expect that, if the Mori program says to replace X by X', then the derived categories of X and X' should be closely related. The most natural language to express such a relation is that of semi-orthogonal decompositions.

A conjecture of Kawamata [Kaw02] gives a precise condition for $\mathrm{D^b}(\mathrm{coh}\, X)$ and $\mathrm{D^b}(\mathrm{coh}\, X')$ to be related by a semi-orthogonal decomposition whenever X and X' are birational.

Definition 3.1.1. Let X and X' be smooth projective varieties. If there are birational morphisms,

$$\begin{array}{ccc} & Z & \\ {\scriptstyle f}\swarrow & & \searrow{\scriptstyle g} \\ X & & X' \end{array}$$

such that $f^*\omega_X \otimes g^*\omega_{X'}^{-1}$ is effective, we say that X **K-dominates** X'.

Conjecture 3.1.2 (Kawamata)**.** *If X K-dominates X', then there is a fully-faithful functor $\mathrm{D^b}(\mathrm{coh}\, X') \to \mathrm{D^b}(\mathrm{coh}\, X)$, hence (automatically by* [BV03]*) a semi-orthogonal decomposition of $\mathrm{D^b}(\mathrm{coh}\, X)$ with $\mathrm{D^b}(\mathrm{coh}\, X')$ as a component.*

As discussed in Section 2.3, the toric Mori program is completely understood. Using this understanding, Kawamata proved Conjecture 3.1.2 in [Kaw05, Kaw06, Kaw12] for smooth projective toric varieties. Moreover, if X' is obtained from X via a step in the toric Mori program, Kawamata exhibited a semi-orthogonal decomposition of $\mathrm{D^b}(\mathrm{coh}\, X)$ with $\mathrm{D^b}(\mathrm{coh}\, X')$ as a component and showed that the complementary component in the semi-orthogonal decomposition possessed a full exceptional collection. In particular,

Theorem 3.1.3 (Kawamata)**.** *Any projective DM toric stack admits a full exceptional collection (not necessarily strong or of line bundles).*

Again, as discussed in Section 2.3, the toric Mori program for a smooth projective toric variety X can be embedded in VGIT by using a presentation of X as a GIT quotient, $\mathbb{A}^n /\!\!/_\chi G$. Hence, the results from [BFK12] on the relationship of derived categories of varieties, or stacks, coming from VGIT can be brought to bear on the problem.

The following result is Theorem 5.2.1 in [BFK12]:

Theorem 3.1.4. *Let X and Y be projective DM toric stacks which are neighbors and let (W, μ) be the associated wall data for a path passing from the chamber for X to the chamber for Y. For any choice of $d \in \mathbb{Z}$ the following statements are true:*

a) *If $\mu > 0$, there exists fully-faithful functors,*

$$\Phi_d : \mathrm{D^b}(\mathrm{coh}\, X) \to \mathrm{D^b}(\mathrm{coh}\, Y)$$

and

$$\Upsilon_- : \mathrm{D^b}(\mathrm{coh}\, W) \to \mathrm{D^b}(\mathrm{coh}\, Y),$$

and a semi-orthogonal decomposition, with respect to Φ_d, Υ_+,

$$\mathrm{D^b}(\operatorname{coh} Y) = \langle \mathrm{D^b}(\operatorname{coh} W)(-\mu - d + 1), \dots, \mathrm{D^b}(\operatorname{coh} W)(-d), \mathrm{D^b}(\operatorname{coh} X)\rangle,$$

with $\mathrm{D^b}(\operatorname{coh} W)(l)$ *the image of* Υ_- *twisted by a character of* λ*-weight* l.

b) *If* $\mu = 0$, *there is an equivalence*

$$\Phi_d : \mathrm{D^b}(\operatorname{coh} X) \to \mathrm{D^b}(\operatorname{coh} Y).$$

c) *If* $\mu < 0$, *there exists fully-faithful functors,*

$$\Phi_d : \mathrm{D^b}(\operatorname{coh} Y) \to \mathrm{D^b}(\operatorname{coh} X)$$

and

$$\Upsilon_+ : \mathrm{D^b}(\operatorname{coh} W) \to \mathrm{D^b}(\operatorname{coh} X),$$

and a semi-orthogonal decomposition, with respect to Φ_d, Υ_+,

$$\mathrm{D^b}(\operatorname{coh} X) = \langle \mathrm{D^b}(\operatorname{coh} W)(-d), \dots, \mathrm{D^b}(\operatorname{coh} W)(\mu - d + 1), \mathrm{D^b}(\operatorname{coh} Y)\rangle,$$

with $\mathrm{D^b}(\operatorname{coh} W)(l)$ *the image of* Υ_+ *twisted by a character of* λ*-weight* l.

Theorem 3.1.4 is the main tool in producing an exceptional collection for any projective toric DM stack, X. See Remark 4.1.16 for a discussion of the meaning of the integer d in the mirror.

The exceptional collection will depend on a choice of presentation,

$$X \cong \mathbb{A}^n /\!\!/_\chi G,$$

of X as a GIT quotient and a path, γ, in the GIT fan for the action of G on $\mathbb{A}^n$. The path, γ, determines a run of the toric Mori program for X.

Theorem 3.1.5. *Let* $X' = \mathbb{A}^n /\!\!/_\chi G$ *be a smooth nef-Fano projective toric DM stack,* γ *be a run of the toric Mori program for* X', *and* X *be a phase along* γ. *Let* $(W_1, \mu_1), \dots, (W_s, \mu_s)$ *be the wall data associated to running* γ *starting at* X. *There exists fully-faithful functors,*

$$\Upsilon_i^j : \mathrm{D^b}(\operatorname{coh} W_i) \to \mathrm{D^b}(\operatorname{coh} X),\ 1 \le j \le \mu_i$$

and a semi-orthogonal decomposition,

$$\mathrm{D^b}(\operatorname{coh} X) = \langle \Upsilon_s^1, \dots, \Upsilon_s^{\mu_s}, \dots, \Upsilon_1^1, \dots, \Upsilon_1^{\mu_1}\rangle.$$

Proof. Let C_0 be the chamber of the GKZ fan on X corresponding to the nef cone of X and C_{-K} be the chamber of the GKZ fan corresponding to the nef cone of X'.

Moving along γ from C_{-K} to C_X, we pass through a finite number of chambers, $C_{-s}, \dots, C_0$, separated by walls, $\tau_{-s}, \dots, \tau_{-1}$, with wall data, $(W_1, \mu_1), \dots, (W_s, \mu_s)$, where $C_{-s} := C_{-K}$ and $C_0 := C_X$. Let X_i be the GIT quotient corresponding to the chamber C_i so that $X_{-s} = X'$ and $X_0 = X$.

Since we are on a run of the toric Mori program, $\mu_i \le 0$ for all i. Therefore by Theorem 3.1.4, there is a semi-orthogonal decomposition,

$$\mathrm{D^b}(\operatorname{coh} X_{i-1}) = \langle \mathrm{D^b}(\operatorname{coh} W_{i-1}), \dots, \mathrm{D^b}(\operatorname{coh} W_{i-1})(\mu_{i-1} - 1), \mathrm{D^b}(\operatorname{coh} X_i)\rangle,$$

where W_{i-1} is the projective toric DM stack appearing in the theorem. Compiling these semi-orthogonal decompositions, we obtain the final desired semi-orthogonal decomposition. □

Let T be an edge-marked tree satisfying the following conditions:

- for each vertex there are at most 2 outgoing edges,
- if a vertex has 2 outgoing edges they are marked by $+$ and $-$,

- if a vertex has 1 outgoing edge it is marked by $+$.

Let X be a toric DM stack. Given a point $p \in \mathcal{F}_{\mathrm{GKZ}}(X)$ define $m(p)$ to be the minimum over the dimension of all cones containing p.

Definition 3.1.6. With notation as above, a **Mori tree** for X is an injective continuous map,

$$\alpha : T \to \mathcal{F}_{\mathrm{GKZ}}(X)$$

satisfying:

- for an edge E marked by $-$ with incoming vertex v, $m(p)$ is constant along $\alpha(E \cup v)$,
- for an edge E marked by $+$ with incoming vertex v and any $p \in E$, $m(\alpha(p)) = m(\alpha(v)) + 1$,
- for any edge E and outgoing vertex v and any $p \in E$, $m(\alpha(p)) = m(\alpha(v)) + 1$
- for any root r of T, $\alpha(r) = 0$.

Corollary 3.1.7. *To any Mori tree α and any GIT presentation of $X \cong \mathbb{A}^n /\!\!/_\chi G$, one can associate an exceptional collection on $\mathrm{D^b}(\mathrm{coh}\, X)$.*

Proof. One can choose a character, χ', in the interior of a chamber of the GIT fan for $\mathbb{A}^n /\!\!/_\chi G$ such that $X' = \mathbb{A}^n /\!\!/_{\chi'} G$ is a nef-Fano projective toric DM stack. We can apply Theorem 3.1.5 to get a semi-orthogonal decomposition

$$\mathrm{D^b}(\mathrm{coh}\, X) = \langle \overbrace{\mathrm{D^b}(\mathrm{coh}\, W_s), \dots, \mathrm{D^b}(\mathrm{coh}\, W_s)}^{-\mu_s}, \dots, \overbrace{\mathrm{D^b}(\mathrm{coh}\, W_1), \dots, \mathrm{D^b}(\mathrm{coh}\, W_1)}^{-\mu_1} \rangle$$

where (W_i, μ_i) are the wall data coming from the line in α given by the $+$ edges. It now suffices to show that each W_i admits a full exceptional collection. From the vertex in α corresponding to the wall for (W_i, μ_i), we can take the straight line path given by first taking the $-$ edge and then all $+$ edges. Again applying Theorem 3.1.5, we have

$$\mathrm{D^b}(\mathrm{coh}\, W_i) = \langle \overbrace{\mathrm{D^b}(\mathrm{coh}\, W'_{s'}), \dots, \mathrm{D^b}(\mathrm{coh}\, W_{s'})}^{-\mu'_{s'}}, \dots, \overbrace{\mathrm{D^b}(\mathrm{coh}\, W'_1), \dots, \mathrm{D^b}(\mathrm{coh}\, W'_1)}^{-\mu'_1} \rangle.$$

where (W'_i, μ'_i) is the wall data associated to this path.

We continuing branching off at $-$ edges to refine the semi-orthogonal decomposition. The process terminates as the dimension of projective toric DM stack strictly decreases each time we branch off using a $-$ edge. The final components are derived categories of zero-dimensional projective toric DM stacks. Such stacks admit full exceptional collections corresponding to all irreducible representations of a finite abelian group. □

Remark 3.1.8. Although the exceptional collection depends on a GIT presentation of $X = \mathbb{A}^n /\!\!/_\chi G$, there is a natural choice: take $\mathbb{A}^n$ to be the spectrum of the Cox ring of X. However, one should note that in the proof above, the GIT fan for W in the wall data (W, μ) is interpreted as the restriction of the GIT fan to the wall. With this in mind, the Cox construction for W as a GIT quotient may have additional torus factors even if X does not.

4. The A-side

4.1. **Degenerations of toric Landau-Ginzburg models.** Let M be as in (2.3). The A-model mirror to a toric nef-Fano stack X is an LG model, i.e., a function

$$w : M \otimes \mathbb{C}^* \to \mathbb{C}. \tag{4.1}$$

The function w is called the **superpotential** and can be obtained as a Laurent polynomial in the following manner ([HV02]).

Write $N = M^\vee = \mathrm{Hom}(M, \mathbb{Z})$ for the dual of M, and consider the dual map from (2.3)

$$\delta^\vee : (\mathbb{Z}^n)^\vee \to N. \tag{4.2}$$

Standard toric descriptions say that X has a stacky fan in N with one-dimensional cones spanned by $\Sigma(1) = \{\alpha_i := \delta^\vee(e_i^\vee) : 1 \leq i \leq n\}$. Let $A = \Sigma(1) \cup \{0\}$ and define z^α to be the character associated to $\alpha \in N$ via the natural isomorphism $N \cong \mathrm{Hom}(M \otimes \mathbb{G}_m, \mathbb{G}_m)$. Then the superpotential is defined as

$$w = \sum_{\alpha \in A} c_\alpha z^\alpha, \tag{4.3}$$

where the coefficients are chosen generically so that all critical points are Morse and have unique critical values.

Remark 4.1.1. As w is defined only using the exact sequence (2.3), toric stacks yielding an identical exact sequence have the same mirror superpotentials. This ambiguity is corrected for upon compactification, as below. As discussed in the introduction, we can only expect that w gives rise to an exact mirror when X is assumed Fano.

Remark 4.1.2. In practice, the set A may be a proper subset of the lattice points of Q. For later constructions it is important that we record this data throughout. Geometrically this data can be encoded in several ways. [GKZ94] introduced a possibly non-normal toric variety X_A whose normalization is the usual projective toric variety X_Q associated to Q. Alternately, A determines a linear subsystem of the ample line bundle $\mathcal{O}_Q(1)$ on X_Q. In [DKK12], the action of the torus on this linear system was used to define a stack $X_{(Q,A)}$. By default this is what is meant by this notation throughout this paper. However, this subtlety will only appear tacitly from here on.

One observes that a choice of generic coefficients of w will not alter the Fukaya-Seidel category $\mathrm{Fuk}^\rightarrow(w)$ as different choices produce equivalent symplectic Lefschetz fibrations (this follows from an application of Moser's trick). This is mirror to the situation on the B-side where the choice of a Kähler structure does not effect the derived category of coherent sheaves. However, if one considers mirror symmetry with the roles of the A- and B-models interchanged (e.g. the Fukaya category of X and the singularity category of w), the coefficients are important and are prescribed by the Kähler structure on X via the mirror map ([CO06, HV02]).

In order to degenerate LG models $w : (\mathbb{C}^*)^n \to \mathbb{C}$, we first compactify the base to $\mathbb{P}^1$ and then partially compactify the total space. To this end, define $X^\vee$ to be the toric stack associated to the convex hull Q of A; this is obtained by considering the normal fan to Q as a stacky fan with all generators primitive. The toric stack $X^\vee$ is equipped with the ample line bundle $\mathcal{O}_Q(1)$ which has a basis of sections given by the lattice points $N \cap Q$. As $A \subset N \cap Q$, we may consider the linear system $V_A \subset H^0(X^\vee, \mathcal{O}_Q(1))$ associated to A, which is simply the span of the monomials $\{z^\alpha : \alpha \in A\}$. We write V_A^{vf} for the torus of all such sections that have non-vanishing coefficients for all $\{z^\alpha\}_{\alpha \in A}$, modulo the diagonal $(1, \ldots, 1)$. We call a section in V_A^{vf} **very full**. Note that by our choice of A we always have a section $s_0 = z^0$ corresponding to the lattice point $\{0\}$. As $X^\vee$ arises from a stacky fan in M, we have that the open dense orbit in $X^\vee$ is $M \otimes \mathbb{G}_m$. The following proposition follows immediately from the definitions.

Proposition 4.1.3. *With notation as above, there exists a unique $s \in V_A^{vf}$ such that the diagram*

$$\begin{array}{ccc} M \otimes \mathbb{G}_m & \longrightarrow & X^\vee \backslash B \\ {\scriptstyle w}\downarrow & & \downarrow{\scriptstyle [s:s_0]} \\ \mathbb{C} & \longrightarrow & \mathbb{P}^1 \end{array} \tag{4.4}$$

commutes, where $B = \mathrm{Zero}(s_0) \cap \mathrm{Zero}(s)$ is the base locus of the pencil $[s : s_0]$. We call $[s : s_0]$ a **mirror pencil** *for X.*

By this proposition, we can equivalently consider the toric mirror superpotential as a one parameter family of hypersurfaces in $X^\vee$ moving in the linear system V_A. One advantage of this perspective is that the moduli theory of such hypersurfaces and their compactifications has been thoroughly studied (e.g [Ale02, DKK12, GKZ94]), and is combinatorial in nature. Note that the torus $T_M = M \otimes \mathbb{C}^*$ acts on V_A^{vf} by coordinate scaling. The following theorem, adapted from Theorem 2.18 of [DKK12], describes the structure of this toric moduli space.

Theorem 4.1.4 ([DKK12])**.** *There exists a toric moduli stack $\mathcal{X}_{\Sigma(A)}$ for very full toric hypersurfaces H and their toric degenerations. In particular, $\mathcal{X}_{\Sigma(A)}$ is a compactification of V_A^{vf}/T_M. The universal hypersurface $\mathcal{H}$ is contained in a toric stack $\mathcal{X}_{\Theta(A)}$ which admits a flat map*

$$\pi : \mathcal{X}_{\Theta(A)} \to \mathcal{X}_{\Sigma(A)}.$$

The polytope $\Sigma(A)$ yielding the stack $\mathcal{X}_{\Sigma(A)}$ is the secondary polytope of (Q, A). The polytope $\Theta(A)$ yielding the stack $\mathcal{X}_{\Theta(A)}$ is a Minkowski sum of $\Sigma(A)$ with a simplex.

The normal fan of the polytope $\Theta(A)$ was introduced by Lafforgue [Laf03] for matroid polytopes. This construction is also implicit in work of Kapranov [Kap95], and the fan appears explicitly in Hacking's subsequent work [Hac03]. The secondary polytope was introduced and thoroughly investigated in [GKZ94], and a more general theory called fiber polytopes was introduced by Billera and Sturmfels [BS92]. This includes as a special case a polytope called the monotone path polytope which we will encounter later. We will not review the full theory of secondary or Lafforgue polytopes here, but will review the most pertinent concepts below.

The most relevant part of this structure for this paper is the 1-skeleton of the secondary polytope. The vertices correspond bijectively to **convex triangulations** T of (Q, A). These are simply cellular decompositions of the polytope Q into lattice simplices whose vertices are respectively contained in the set A, and subject to the condition that they arise from the projection of the linearity domains of a piecewise affine convex function. Given such a T, the corresponding vertex $\phi_T \in \mathbb{R}^A$ of $\Sigma(A)$ is given simply by

$$\phi_T^\alpha = \sum_{\alpha \in \sigma} \mathrm{vol}(\sigma) \tag{4.5}$$

where ϕ_T^α denotes the corresponding coordinate in $\mathbb{R}^A$, and the sum is over the normalized lattice volumes of all simplices containing the lattice point $\alpha \in A$. Of particular interest to us are the edges of $\Sigma(A)$ which correspond bijectively to extended circuits of (Q, A).

Definition 4.1.5. An (affine) **circuit** in a lattice L is a set $C \subset L$ which is affinely dependent over L, but such that every proper subset is affinely independent. If L has rank d, an **extended circuit** is any subset $\tilde{C} \subset L$ such that $|\tilde{C}| = d + 2$ and which affinely spans L.

It is easy to check that any extended circuit contains a unique circuit. We may write an extended circuit explicitly by writing the affine relation

$$\sum_{\alpha\in A} c_\alpha \alpha = 0$$

where $\sum_{\alpha\in A} a_\alpha = 0$. Let $A^+ = \{\alpha \,|\, a_\alpha > 0\}$, $A^- = \{\alpha \,|\, a_\alpha < 0\}$, and $A^\circ = \{\alpha \,|\, a_\alpha = 0\}$. The triple $(|A^+|, |A^-|; |A^\circ|)$ is an invariant of an extended circuit called the **signature**, and is well-defined up to a switch of the first two entries. If $|A^\circ| = 0$, we will write $(|A^+|, |A^-|)$ for the signature and refer to the extended circuit as non-degenerate. The relationship between (extended) circuits and triangulations is that every extended circuit determines two (convex) triangulations $T^\pm$ of the polytope $Q = \mathrm{conv}(A)$ by taking

$$T^\pm = \{\mathrm{conv}(A \setminus \{\alpha_i\} \,|\, \alpha_i \in A^\pm\}$$

as the decomposition of Q into simplices with vertices contained in A.

Remark 4.1.6. A similar theory can be developed with simplicial sub-fans supported on a set of rays playing the role of triangulations of a marked polytope. Here a (linear) circuit will correspond to sets of primitive generators which are linearly dependent but such that every proper subset is linearly independent, and these linear circuits will correspond bijectively to the walls of the GKZ fan. Indeed, the entire theory of the GKZ fan of a toric variety can be developed in this language; this was the viewpoint taken in the original work of Reid [Rei83] and Oda-Park [OP91], and is discussed in detail in [CLS11, Section 15.3].

When X is a toric Fano variety, the relationship between the GKZ fan and the secondary polytope is particularly clear. Let Δ be a reflexive polytope defining X, Δ° to be the polar dual, and define the set $A = \mathrm{vert}(\Delta^\circ) \cup \{0\}$. The following result is well-known, see e.g. [Cox98, Section 7].

Proposition 4.1.7. *With notation as above, the enlarged GKZ fan from Proposition 2.3.12 equals the normal fan of the secondary polytope $\Sigma(A)$. In particular, the GKZ fan of X is a full-dimensional subfan of the normal fan of $\Sigma(A)$.*

This proposition will be used at least implicitly many times in order to identify walls in the GKZ fan (B-side) with circuits on the mirror (A-side).

Let us now interpret the above combinatorial constructions geometrically. Following [Ale02], one associates to T (or more generally any face of $\Sigma(A)$) a **stable toric variety** which is a possibly reducible variety whose components are the toric stacks corresponding to the respective cell of the subdivision of Q, and which are glued together along boundary strata according to the combinatorics of the subdivision. A fiber of $\mathcal{H} \to \mathcal{X}_{\Sigma(A)}$ is a **degenerated hypersurface**; namely a collection of reducible hypersurfaces in each of the components of a stable toric variety with the respective gluing. The structure of a degenerated hypersurface is closely related to the corresponding tropical variety; we review this in 5.1.1.

A specific consequence of the above theorem is that one can realize a mirror pencil from 4.1.3 as the pullback along a map $\iota : \mathbb{P}^1 \to \mathcal{X}_{\Sigma(A)}$ as in the diagram below

$$\begin{array}{ccccc} M \otimes \mathbb{G}_m & \xrightarrow{i} & X^\circ \backslash B & \xrightarrow{\iota^*} & \mathcal{H} \\ {\scriptstyle \mathbf{w}}\downarrow & & {\scriptstyle [s:s_0]}\downarrow & & {\scriptstyle \pi|_{\mathcal{H}}}\downarrow \\ \mathbb{C} & \longrightarrow & \mathbb{P}^1 & \xrightarrow{\iota} & \mathcal{X}_{\Sigma(A)} \end{array} \tag{4.6}$$

We use this description to construct the moduli space of LG models; that is, the moduli space of mirror pencils.

Remark 4.1.8. The theory of maps from a sphere to a symplectic variety is quite rich and, were $\mathcal{X}_{\Sigma(A)}$ a smooth variety (or DM stack), we could study the structure of the (twisted) stable map compactification $\mathcal{M}_0(\mathcal{X}_{\Sigma(A)}, \iota(\mathbb{P}^1))$ in order to give a useful description of the space of all such mirror pencils. Unfortunately, it is quite rare that $\mathcal{X}_{\Sigma(A)}$ is smooth even as a stack. The approach we give instead better reflects the underlying combinatorics.

Let

$$\rho : V_A^{vf} \to \mathcal{X}_{\Sigma(A)} \tag{4.7}$$

be the map which is the composition of the quotient $V_A^{vf} \to V_A^{vf}/T_M$ with the inclusion of V_A^{vf}/T_M onto the dense torus of $\mathcal{X}_{\Sigma(A)}$. Now, let $K = \mathbb{G}_m$ be the one parameter subgroup of the open torus of $\mathcal{X}_{\Sigma(A)}$ which acts via

$$\lambda \cdot \rho\left(\sum_{\alpha\in A} c_\alpha s_\alpha\right) = \rho\left(\lambda c_0 s_0 + \sum_{0\neq\alpha\in A} c_\alpha s_\alpha\right). \tag{4.8}$$

By varying $s = \sum_{\alpha\in A} c_\alpha s_\alpha$ and comparing with the definition of a mirror pencil as in Proposition 4.1.3, we obtain the following useful characterization.

Proposition 4.1.9. *Any mirror pencil for X is represented by the closure of a K-orbit in $\mathcal{X}_{\Sigma(A)}$.*

Being a one-parameter subgroup, K corresponds equivalently to a cocharacter; let us now describe it explicitly. Consider the exact sequence

$$0 \to M \oplus \mathbb{Z} \xrightarrow{\tilde{\delta}} \mathbb{Z}^A \xrightarrow{\tilde{\gamma}} \widehat{G} \to 0$$

where $\tilde{\delta}(m, \lambda) = \sum_{\alpha\in A}(\langle m, \alpha\rangle + \lambda)$, and $\widehat{G}$ is defined to be the cokernel. This exact sequence is just the A-side analogue of (2.3). Recall that in our situation $A = \mathrm{vert}(\Delta) \cup \{0\}$, and let $A' = \mathrm{vert}(\Delta)$, which we may identity with the primitive generators of the fan defining the original toric Fano.

Comparing with the exact sequence (2.3), consider the commutative diagram

$$\begin{array}{ccccccccc}
0 & \longrightarrow & M & \xrightarrow{\delta} & \mathbb{Z}^{A'} & \xrightarrow{\gamma} & \widehat{G} & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle i_1} & & \downarrow{\scriptstyle i_2} & & \downarrow{\scriptstyle =} & & \\
0 & \longrightarrow & M \oplus \mathbb{Z} & \xrightarrow{\tilde{\delta}} & \mathbb{Z}^A & \xrightarrow{\tilde{\gamma}} & \widehat{G} & \longrightarrow &
\end{array}$$

where i_1 is inclusion in the first factor, i_2 is the inclusion $\mathbb{Z}^{A'} \subset \mathbb{Z}$, and

$$\tilde{\delta}(m, k) = i_2(\delta(m)) + k\sum_{\alpha\in A} e_\alpha. \tag{4.9}$$

The cocharacter of K acting on $\mathbb{Z}^A \otimes \mathbb{G}_m$ (before applying ρ) is clearly e_0. This descends via $\tilde{\gamma}$ to the cocharacter $\chi = \tilde{\gamma}(e_0)$.

Remark 4.1.10. Note that as an element of $\widehat{G}$, the cocharacter χ corresponding to the subgroup K coincides with the anticanonical divisor of the original variety X.

It is thus natural to define the space of mirror pencils to be a quotient of the stack $\mathcal{X}_{\Sigma(A)}$ by K. This requires the correct notion of quotient. The most structured approach is to use the exact sequences above, along with the generalized notion of a toric stack developed in [GS11], to take this quotient at the level of generalized stacky fans. If we only regard the toric pencils as equivariant cycle classes determined by $\iota(\mathbb{P}^1)$, we may take a more classical approach.

Definition 4.1.11. The space of LG mirrors X^{LG} is the toric Chow quotient of $X_{\Sigma(A)}$ by K.

For the definition and properties of this quotient, see [KSZ91]. Note that in general this will be a non-normal toric variety. It was observed in [CS12] that this space admits an equivalent description in terms of log stable maps. It is important to note that the polytope corresponding to X^{LG} was discussed as a purely combinatorial object in [BS92], where it was dubbed the **monotone path polytope**. More precisely, this polytope is the monotone path polytope of $\Sigma(A)$ relative to the hyperplane. The most important feature of this polytope is that its vertices correspond to edge paths on the secondary polytope. Intuitively, the cocharacter $\chi = \tilde{\gamma}(e_0)$ determines a linear functional on the space $\widehat{G}_{\mathbb{R}}$ which contains the secondary polytope, and the edge paths are those which move "downward" with respect to this linear functional.

Taking a page from mirror symmetry of Calabi-Yau varieties, we consider Homological Mirror Symmetry of the original nef-Fano stack as taking place near the toric fixed points in X^{LG}; that is, near the large complex structure limit.

Definition 4.1.12. A torus fixed point of X^{LG} is called a maximally degenerated LG mirror.

The following theorem relates these degenerated LG models to their B-side mirrors, and makes precise the notion that the edge paths specified by vertices of the monotone path polytope are mirror to runs of the Mori program. A complete proof of this can be found in [DKK12],

Theorem 4.1.13 ([DKK12])**.** *Maximal degenerated LG mirrors are in bijection with straight line run of the toric Mori program on X.*

Let us now describe the geometry of maximally degenerated LG mirrors in more detail. As mentioned above, they in particular correspond to a collection of edge paths on the secondary polytope, and thus a sequence of circuits. Geometrically, to every maximal degeneration $\psi \in X^{LG}$ there is thus an associated 1-cycle

$$Z_\psi = \cup_{i=1}^m a_i Z_{\psi,i} \tag{4.10}$$

on $\mathcal{X}_{\Sigma(A)}$. Equivalently, this is a degeneration of the generic mirror pencil 4.3 into a (weighted) union of one dimensional toric boundary strata. The coefficients of the cycle can be seen by taking $\{\psi(t)\}_{t\in[0,1]} \subset X^{LG}$ to be a one dimensional family of LG pencils with $\psi(0) = \psi$ a maximal degeneration, and observing a bubbling phenomena for the corresponding maps $\psi(t) : \mathbb{P}^1 \to \mathcal{X}_{\Sigma(A)}$ into a_i-ramified maps

$$\psi_i(0) : \mathbb{P}^1 \to Z_{\psi,i}. \tag{4.11}$$

By the description of the universal hypersurface $\mathcal{H}$ over $\mathrm{X}_{\Sigma(A)}$, we thus see that ψ defines a chain of (stacky) $\mathbb{P}^1$'s, which is equipped with a family of degenerated hypersurfaces. Over the nodes of this chain and over the two torus fixed points at the end of the chain the

hypersurface is typically reducible and has irreducible components consisting of (stacky) pairs of pants glued according to the corresponding triangulation. The generic fiber over a components of the chain will likewise be determined by the circuit determined by the two triangulations of the respective endpoints. The advantage of these degenerate pencils is that they are essentially entirely combinatorial. In practice, we will need Lefschetz pencils which approximate these degenerate pencils. This can be done by considering a point in X^{LG} sufficiently close to the corresponding fixed point. We will call such a pencil a **regeneration** of ψ. In practice, there are subtleties due to the non-trivial stack structure of $\mathcal{X}_{\Sigma(A)}$, see Section 4.3 of [DKK12] for details.

There are prescribed asymptotics in t for annuli B_i in the domain of $\psi(t)$ for which $\psi(t)|_{B_i}$ converges to the components $\psi_i(t)$ ([DKK12]). These asymptotics separate the critical values of $\psi(t)$ into subsets $\{S_1, \ldots, S_m\}$. Furthermore, as $\psi_i(t)$ converges to an a_i-fold cover of Z_i, each subset S_i of critical values further partitions into a_i groups $\{S_{i1}, \ldots, S_{ia_i}\}$ corresponding to the branches of the cover. To summarize, given a maximal degeneration $\psi(0) = \psi \in X^{LG}$, and a one-parameter family of LG-models approaching it, we obtain a decomposition of the plane into successive annuli such that:

- There is one annulus for each circuit appearing in the edge path on $\Sigma(A)$ determined by ψ
- If an edge appears with multiplicity, the corresponding annulus is divided into the corresponding number of radial regions.

The number of critical points in each such region then depends on the further geometry of the corresponding circuit. In particular, it is 1 unless the circuit is what is known as a degenerate circuit. To compute the multiplicities a_i, let $\mathrm{Stab}(Z_i)$ be the subtorus in $\widehat{G} \otimes \mathbb{C}^*$ that stabilizes Z_i. Then a_i equals the order of the group $\mathrm{Ker}(\eta_i)$ where

$$\eta_i : K \to \widehat{G}/\mathrm{Stab}(Z_i). \tag{4.12}$$

The computation of this order amounts to evaluating the cocharacter of K with a generator for the affine edge $\{T_i, T_{i+1}\}$ in the secondary polytope $\Sigma(A)$ corresponding to Z_i. In particular, if φ_{T_i} and $\varphi_{T_{i+1}}$ are the vertices on the edge $\Sigma(A)$ regarded as a polytope in $\mathbb{Z}^A$ and their difference $\varphi_{T_{i+1}} - \varphi_{T_i}$ is not a nontrivial multiple of any lattice point, then

$$a_i = \left\langle e_0, \varphi_{T_{i+1}} - \varphi_{T_i} \right\rangle \tag{4.13}$$

The total collection of the decomposition into annuli and the distinguished basis is called a **radar screen**. We will see an explicit example in Figure 14.

Remark 4.1.14. For a run of the Mori program, let (W_i, μ_i) be the wall data associated to an affine edge $\{T_i, T_{i+1}\}$. It is shown in [DKK12] that

$$a_i = -\mu_i \cdot \dim \mathrm{K}_0(W_i).$$

When the corresponding extended circuit is non-degenerate $a_i = -\mu_i$. Notice that there's a minus sign due to an altercation among sign conventions in [DKK12] and [BFK12]. In addition, the locus W is determined by the locus where the fan is changed nontrivially i.e. the support of the circuit (see [CLS11, Section 15.3]).

We may use this result to write down a conjectural mirror for the semi-orthogonal decomposition given in Theorem 3.1.5. Strictly speaking, if X' is a nef-Fano and X is a phase, the theorem below gives the conjectural mirror semi-orthogonal decomposition of $\mathrm{D^b}(\mathrm{coh}\, X)$, of

which a subset of components specified by omitting the wall contributions from X to X' yields $\mathrm{D}^{\mathrm{b}}(\operatorname{coh} X')$.

Theorem 4.1.15 ([DKK12])**.** *For every maximal degenerated LG mirror ψ of a family $\psi(t)$ there is a semi-orthogonal decomposition*

$$\langle \overbrace{\mathrm{Fuk}^{\rightarrow}(\psi_1),\ldots,\mathrm{Fuk}^{\rightarrow}(\psi_1)}^{-\mu_1},\ldots,\overbrace{\mathrm{Fuk}^{\rightarrow}(\psi_s),\ldots,\mathrm{Fuk}^{\rightarrow}(\psi_s)}^{-\mu_s}\rangle.$$

of the Fukaya-Seidel category $\mathrm{Fuk}^{\rightarrow}(\psi(t))$ for $t \approx 0$. Here the ψ_i are regenerations of the singular pencils Z_i.

Remark 4.1.16. The integer d from Theorem 3.1.4 is mirror to a monodromy transformation around a degenerate fiber of Z_i (i.e. a fiber over a torus fixed point of the base stacky $\mathbb{P}^1$), where Z_i is the pencil corresponding to the circuit which is mirror to the corresponding wall crossing. In terms of the radar screen, this amounts to a rotation of the corresponding annulus and hence yields a different semi-orthogonal decomposition of the Fukaya-Seidel category $\mathrm{Fuk}^{\rightarrow}(\psi(t))$ for $t \approx 0$.

5. The Mori program meets Homological Mirror Symmetry, torically

5.1. **General method.** We begin this section by recalling our central conjecture 5.1.1. Following the notation of the previous sections, we let X be a projective toric DM stack, X' be a projective toric nef-Fano DM stack with a straight line run for X containing X as a phase. We take $\psi = \psi(0)$ to be the degenerated LG mirror corresponding to the Mori program run and $\{\psi(t)\}$ a degenerating family.

Conjecture 5.1.1. *Given any maximal degeneration ψ of an LG mirror pencil, let*

$$\begin{aligned} \mathrm{Fuk}^{\rightarrow}(\psi(t)) &= \langle \mathcal{T}_1,\ldots,\mathcal{T}_r\rangle, \\ \mathrm{D}^{\mathrm{b}}(\operatorname{coh} X) &= \langle \mathcal{S}_1,\ldots,\mathcal{S}_r\rangle \end{aligned}$$

be the semi-orthogonal decompositions associated to ψ and its run of the toric Mori program.

Then there exists an equivalence of triangulated categories

$$\Phi_\psi : \mathrm{Fuk}^{\rightarrow}(\psi(t)) \to \mathrm{D}^{\mathrm{b}}(\operatorname{coh} X)$$

which restricts to equivalences $\Psi_\psi : \mathcal{T}_i \to \mathcal{S}_i$ for all $1 \le i \le r$.

In addition to being a statement of Homological Mirror Symmetry for toric varieties, the conjecture is formulated to suggest a rich correspondence between decompositions of the categories involved, the geometry of the space of LG models, and birational geometry.

Before outlining a strategy to prove this conjecture, we cite some evidence in its favor. It was proven in [DKK12] that the conjecture holds at the level of K-theory ranks.

Proposition 5.1.2 ([DKK12],5.17)**.** *In the decompositions of Conjecture 5.1.1*

$$\operatorname{rank}(K_0(\mathcal{T}_i)) = \operatorname{rank}(K_0(\mathcal{S}_i)) \tag{5.1}$$

for all $1 \le i \le r$.

There are two hurdles that must be overcome in order to prove Conjecture 5.1.1. The first is to show that

$$\mathcal{T}_i \simeq \mathcal{S}_i \tag{5.2}$$

for any circuit in $\Sigma(A)$, or equivalently, for any wall crossing in the GKZ-fan coming from a straight line run. The second hurdle is showing that the A_∞- or dg-bimodule of morphisms from $\langle \mathcal{T}_1, \dots, \mathcal{T}_k \rangle$ to $\mathcal{T}_{k+1}$ is quasi-equivalent to that from $\langle \mathcal{S}_1, \dots, \mathcal{S}_k \rangle$ to $\mathcal{S}_{k+1}$. The full equivalence would then follow from an easy induction. For both the A and B side of mirror symmetry, a detailed study of the dg- or A_∞- endomorphism algebra associated to exceptional collections in the respective category must be made.

5.1.1. *Calculating the A_∞-algebra corresponding to a run of the toric Mori program on the A-side.* For the A-side, we sketch a correspondence between LG models and a type of "tropical Morse theory". The main goal of which is to understand the structure of the Lagrangian vanishing cycles which appear in the regenerated LG models, as this is precisely what determines $\mathrm{Fuk}^{\rightarrow}(\psi)$ by Theorem 4.1.15. The main point is that the location of these vanishing cycles can be largely described by determining their tropicalizations. The idea is simple: recall that for each component in the chain of (stacky) $\mathbb{P}^1$'s given a maximal LG degeneration ψ its two torus fixed points correspond to triangulations, and the entire components corresponds to a circuit. We will show that the tropicalization of the vanishing cycles of a regeneration are determined by the circuit data.

To this end, we first recall some basic notions from tropical geometry. First, recall that given a polarized toric variety X_Q determined by a lattice polytope Q, there is a continuous map

$$\mathrm{Mo} : X_Q \to Q$$

called the moment map, such that the fiber over a point in the interior of a codimension k cell of Q is the real torus $(S^1)^k$. In practice, it will suffice to consider the restriction of Mo to the dense torus, in which case, identifying the interior of Δ homeomorphically with $\mathbb{R}^n$, the map is identified with the usual amoeba map

$$\mathrm{Log}(t_1, \dots, t_n) = (\log|t_1|, \dots, \log|t_n|).$$

If $H \subset X_Q$ is a hypersurface, the image $\mathrm{Mo}(H) \subset Q$ is called the **amoeba** of H. Tropical geometry studies the structure of H via polyhedral complexes onto which the amoeba deformation retracts.

More precisely, given a convex triangulation T of a polytope (Q, A), we may define a tropical variety Y_T which is the dual polyhedral complex to the triangulation, with maximal cells weighted by the lattice volume of the corresponding dual simplex. To each simplex in the triangulation, we associate a weighted tropical pair of pants which glue together to give the full tropical variety Y_T.

Recall from (4.5) that T corresponds to a vertex of the secondary polytope $\Sigma(A)$ and thus a unique fixed point p_T of the stack $\mathcal{X}_{\Sigma(A)}$. The relationship between the geometry of a hypersurface and Y_T is most clear when the triangulation is assumed to be **maximal**; that is, when the maximal simplices of T each have normalized lattice volume equal to 1. As before, if $p \in \mathcal{X}_{\Sigma(A)}$, we let $\mathcal{H}_q$ denote the hypersurface in X_Q which is the fiber of $\pi_{\mathcal{H}}$ over p. Let $\mathcal{H}_p^\circ$ denote the open part of this hypersurface which is contained in the dense torus of X_Q.

The following proposition follows immediately from the main results of [Mik04].

Theorem 5.1.3. *Let T be a maximal triangulation of (Q, A), and let $q(t)$ be a continuous path in $\mathcal{X}_{\Sigma(A)}$ such that $q(0) = p_T$ and $q(t)$ does not meet the toric boundary of $\mathcal{X}_{\Sigma(A)}$ for $0 < t < \epsilon$. Then as $t \to 0$, the amoebas $\mathrm{Mo}(\mathcal{H}_{q_t}^\circ)$ converge, in the Gromov-Hausdorf limit, to the tropical variety Y_T.*

Analyzing the convergence for sufficiently small t produces a continuous map

$$\pi^{trop} : \mathcal{H}^{\circ}_{q_t} \to Y_T \tag{5.3}$$

called a **tropical degeneration** of $\mathcal{H}_{q_t}$. The fibers of this map over the interior of a maximal cell are simply real tori.

We now apply this machinery to the study the fibers of mirror pencils. Recall from Section 4 that a maximal degeneration of a mirror pencil arises as the limit of a family $\{\phi(t)\}$ of mirror pencils whose limit in the space X^{LG} determines a sequence of edge paths $\{T_i, T_{i+1}\}$. Thus, for each $1 \leq i \leq m$ and $t \in [0,1]$ we may define $\{q_{i,t}\}$ on the boundary of the i-th annulus B_i of in the K-orbit $\phi(t)$ which converges to p_{T_i}. In particular, the fibers near the maximal degeneration ϕ can be viewed as going through a sequence of tropical degenerations

$$\pi_i^{trop} : \phi(t)^*(Z_{q_{i,t}}) \to Y_{T_i}. \tag{5.4}$$

We now discuss how to use tropical techniques to study not just the fibers over the points $q_{i,t}$, but the vanishing cycles which appear in the regenerated components ψ_i from Theorem 4.1.15. It was noted in [GKZ94, Example 11.5.8] that there is a strong relationship between triangulations related by a circuit and Morse surgery. This motivates the following definition.



FIGURE 6. Unstable and stable complexes for two dimensional circuits.

Definition 5.1.4. Let T and T' be triangulations related by a circuit, and let Y_T and $Y_{T'}$ be the respective tropical varieties. A cell $P \subset Y_{T'}$ is a **stable cell** with respect to the circuit if its dual simplex is contained in T' but not T. The **unstable cells** are those in Y_T whose dual simplices are contained n T but not T'. The collection of all stable cells will be called the **stable complex** of the modification.

Of course, the definition depends on an ordering of $\{T, T'\}$. In all situations we consider, this is afforded by the sequence $T_0, \ldots, T_n$ defined by a maximal LG degeneration ψ. In Figure 6 we illustrate the stable cells in red and the unstable cells in blue for non-degenerate circuits in two dimensions. The main reason for these considerations is to apply the technology of stable and unstable complexes to the study of vanishing cycles which appear in degenerations of LG models. We prove here that the stable complex is indeed the tropicalization of a vanishing cycle.

Proposition 5.1.5. *Let $\{\psi(t)\}$ be a maximal degeneration of a mirror potential with $\{T_0, \dots, T_m\}$ the corresponding sequence of triangulations. If γ is a path in B_i from a critical value of $\psi(t)$ to $q_{i,t}$ with vanishing cycle L, then the image $\pi_i^{trop} : L \to Y_{T_i}$ converges to the stable complex of $\{T_i, T_{i+1}\}$ as $t \to 0$.*

Proof. Recalling Theorem 4.1.4, we have the map $\pi : \mathcal{X}_{\Theta(A)} \to \mathcal{X}_{\Sigma(A)}$ along with its restriction $\pi_{\mathcal{H}}$ to the universal hypersurface. On the annulus B_i, the maps $\psi(t) : \mathbb{P}^1 \to \mathcal{X}_{\Sigma(A)}$ converge to an a_i-fold cover of the $\mathbb{G}_m$ orbit corresponding to the edge $\{T_{i-1}, T_i\}$ on the polytope $\Sigma(A)$. Let S be the convex subdivision corresponding to the edge $\{T_{i-1}, T_i\}$ and note that the polytopes in S consist of the simplices that are members of both triangulations and the polytope C which contains all simplices of which are being modified.

The pullback via $\psi(0)$ of the hypersurface $\mathcal{H} \subset \mathcal{X}_{\Theta(A)}$ along the component Z_i decomposes into components

$$\psi(0)|^*_{Z_{\psi,i}}(\mathcal{H}) = \cup_{\sigma \in S} X_\sigma \tag{5.5}$$

where X_σ is a fiber bundle of a of pants (or a cover thereof) over $\mathbb{P}^1$ for all $\sigma \neq C$ and the map $f_C : X_C \to \mathbb{P}^1$ is associated to the subpolytope $C \subset Q$ associated to the circuit. It is a consequence of Proposition 5.7 of [DKK12] that the critical points of $\psi(t)|_{B_i}$ converges to the unique critical point of f_C and is contained in the interior of X_C.

Let $\tilde{Y}_{T_i}$ be the tropical variety associated to the T_i triangulation of the marked polytope $(C, C \cap A)$. Then, there is an inclusion of the polyhedra in $\tilde{Y}_{T_i}$ to those in Y_{T_i}. We observe that at $t = 0$, the tropical image of the vanishing cycle L is contained dual complex $\tilde{Y}_{T_i} \subset Y_{T_i}$, as it is the same image as that of the hypersurfaces corresponding to f_C. One can use a compressing flow to force this image to be in the compact polyhedra of $\tilde{Y}_{T_i}$. It is an easy exercise to see that for a circuit, these polyhedra give the stable complex of the circuit corresponding to $\{T_i|_C, T_{i+1}|_C\}$. But it easily follows from the definition of stable complex that the inclusion of the stable complex in $\tilde{Y}_{T_i}$ of $\{T_i|_C, T_{i+1}|_C\}$ into the full tropical complex Y_{T_i} will be a bijection of polyhedra. This yields the claim. □

Remark 5.1.6. Conjectures of Seidel [Sei11] suggest that it may be possible to build a calculus that yields the Floer theory of the vanishing cylces in terms of essentially tropical data. The above proposition indicates that one can identify the vanishing cycles of an LG mirror with cell complexes in the tropical hypersurface Y_{T_m} near infinity; however, the results in [Abo09] strongly suggest that, in general, in order to fully recover the Floer theory of a Lagrangian from its tropicalization, one requires additional combinatorial enhancements. In addition to standard tropical techniques, establishing related results would require the study of Floer theory in the coamoeba setting [FHKV09, She11]. Were such a calculus to be built, the A_∞-endomorphism algebra of the exceptional collection of vanishing cycles would then be read off from the combinatorial data of the tropical vanishing cycles, after which a comparison with the B-model could be made. The implementation of this program is beyond the scope of the work, but motivates much of our analysis. In particular, we view the analysis of the A-side of the main example in Section 5.2 as a particular instance of this philosophy.

In the case where the fibers are Riemann surfaces, using matching paths arguments, the actual vanishing cycles can be explicitly identified for the non-degenerate circuits (of which there are only two cases). These cycles are pictured in Figures 6 and 7 as an illustration of Proposition 5.1.5.



FIGURE 7. Circuit vanishing cycles

In the three cases on the right of Figure 7, the vanishing cycle is embedded in the hypersurface while in the case on the left, one sees an immersed vanishing cycle which is a $(\mathbb{Z}/3\mathbb{Z})$ quotient of the hexagonal torus. In this last case, the actual Lagrangian lies in the cover. The monodromy around the fixed points of these circuit fibrations is given by a Dehn twist along each cylinder that corresponds to an edge in the stable or unstable manifold. In the case of the finite quotient, this is simply the action of $(\mathbb{Z}/3\mathbb{Z})$ on the cover. In the next section, we pull these cycles back along the tropical hypersurfaces associated to a maximal LG mirror degeneration.

Remark 5.1.7. Compared to computing Floer cohomology, there are a multitude of general ways for computing morphisms in the derived category of coherent sheaves on a variety. However, the A-model analysis described above and the methods and observations of [AS10, Sei11] suggest an approach to the B-side computations of A_∞ structure that might be amenable to mirror symmetry. Namely, the hypersurface on the A-side admits a pair pants decomposition coming from its tropical amoeba. This decomposition is combinatorially dual to the collection of maximal cones in the fan for the corresponding toric variety. If we wish to compute morphisms on the A-side, following [Sei11], we should be able to compute the Floer cohomology of the restrictions of the Lagrangians to each pair of pants and glue them together to get the global Floer cohomology. Mirror to this procedure, we should use the affine cover of the toric variety coming from the maximal cones, plus locally-free resolutions of the complexes mirror to the Lagrangians. Hence, we obtain a dg-algebra from the corresponding Čech resolution. However, due to the fact that much of this technology is not currently available, this is not the approach we will take in Section 5.2.

5.2. **An example.** To illustrate the program outlined above, we examine the case of a blow up of $\mathbb{P}^2$ in three infinitesimally close points. Precisely, take $\mathbb{P}^2$, blow it up at a point, and then blow up at the two points on the exceptional locus corresponding to two coordinate axes passing through the original point. This is toric. Namely, we take X to be the toric variety defined by the fan Σ in $N_{\mathbb{R}}$ with one cone generators

$$\Sigma(1) = \{(-1,-1)(1,0),(0,1),(1,1),(1,2),(2,1)\}. \tag{5.6}$$

The fan Σ is pictured in Figure 8 with rays labeled according to the ordering in the above equation. We name the corresponding toric divisors $\{D_0,\dots D_5\}$.

This fan determines the linear map γ from $\mathbb{Z}^6$ to $\widehat{G} \cong \mathbb{Z}^4$ displayed in Figure 9 with basis for $\mathbb{Z}^4$ given by D_0, D_3, D_4, D_5.

The variety X is not nef-Fano. However, if $\Sigma'(1) = \Sigma(1)\backslash\{(1,1)\}$ and X' is the smooth stack associated to Σ', one observes that X' is nef-Fano and X' is a phase of X. Figure 10



FIGURE 8. The fan of the toric variety X

$$\gamma = \begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 1 & 0 & 0 \\ 0 & -2 & -1 & 0 & 1 & 0 \\ 0 & -1 & -2 & 0 & 0 & 1 \end{pmatrix}$$

$$\mathbb{Z}^6 \longrightarrow \mathbb{Z}^4 \cong \widehat{G}$$

FIGURE 9. The map γ for the toric variety X

illustrates a straight line run of the toric Mori program for X' with Σ' appearing on the far left and Σ appearing as the first phase. The one-parameter subgroups determining the wall-crossings are listed above each fan modification in the basis for D_0, D_3, D_4, D_5.



FIGURE 10. A straight line run of the toric Mori program for X' marked by one-parameter subgroups

Following the outline given in Section 4, the straight line run of the toric Mori program appearing in Figure 10 can also be interpreted in the language of monotone path polytopes. To this end, consider the secondary polytope of $A = \Sigma(1) \cup \{(0,0)\}$. Recall that the normal fan of the polytope extends the GIT fan of X. A direct calculation finds that the secondary polytope $\Sigma(A)$ is a 4-dimensional polytope with 30 vertices, 20 of which have normal cones in the GIT fan of X. The monotone path polytope of $\Sigma(A)$ with respect to the function χ can also be computed as a 3-dimensional polytope with 24 vertices. In this case, every run

of the toric Mori program for X' is a straight line run and so the vertices of the monotone path polytope are in one to one correspondence with straight line runs of the toric Mori program of X'. Of these runs, precisely 10 have X as a phase. The monotone path polytope is illustrated in Figure 11. Those faces which have vertices whose runs of the toric Mori program contain X as a phase are colored in green. The left most green vertex corresponds to the straight line run of the toric Mori program illustrated in Figure 10.



FIGURE 11. The monotone path polytope $\Sigma_\chi(\Sigma(A))$.

5.2.1. *The B-model computation.* Let us now discuss the dg-algebra associated to this straight line run of the toric Mori program on the B-side.

Recall that X is, by definition, the toric variety defined by the fan Σ in $N_{\mathbb{R}}$ with one cone generators labeled from 0 to 5 by

$$\Sigma(1) = \{(1,0),(2,1),(1,1),(1,2),(0,1),(-1,-1)\}. \tag{5.7}$$

The Cox construction describes X as a GIT quotient of the prime spectrum of the homogeneous coordinate ring, $\mathbb{C}[x_0,\ldots,x_5]$, by the torus dual to the Picard group of X,

$$G := \mathrm{Hom}(\mathrm{Pic}(X), \mathbb{G}_m),$$

for any choice of a G-equivariant ample line bundle. In this case, the semi-stable locus and stable locus agree. They are the complement of the vanishing locus of the ideal

$$J := \langle x_3x_4x_2x_0, x_4x_2x_0x_1, x_2x_0x_1x_5, x_0x_1x_5x_3, x_1x_5x_3x_4, x_5x_3x_4x_2\rangle.$$

Let

$$U := \mathrm{Spec}\,\mathbb{C}[x_0,\ldots,x_5]\backslash V(J)$$

so that

$$X \cong [U/G]. \tag{5.8}$$

Via the above isomorphism, one can identify D_i with the vanishing locus of x_i for $0 \leq i \leq 5$. With this interpretation, the two disjoint exceptional loci from the second blow-up are D_4 and D_5 while the strict transform of the exceptional locus from the first blow-up is described by D_3. The remaining toric divisors, D_0, D_1 and D_2 are all linearly equivalent to the pullback of a generic hyperplane from $\mathbb{P}^2$.

Now, the run of the Mori program appearing in Figure 10 (starting from the ample chamber of X and ending outside the support of the GKZ fan) corresponds to the sequence of blow-downs for X given by D_2 followed by D_4 followed by D_3. Consider the blow down maps

$$f_5 : X \to Y$$

contracting D_5,

$$f_4 : Y \to \mathbb{F}_1$$

contracting D_4, and

$$g : \mathbb{F}_1 \to \mathbb{P}^2$$

contracting D_3.

Table 1 displays the weights of each of the one-parameter subgroups appearing in Figure 10 along with the data (W, μ) used to determine the semi-orthogonal decomposition in Theorem 3.1.5. The one-parameter subgroup λ_1 determines the variation of GIT quotients from X' to X. Therefore, to compute the semi-orthogonal decomposition of $\mathrm{D}^{\mathrm{b}}(\mathrm{coh}\, X)$ we consider the sequence of wall-crossings corresponding to $\lambda_2, \lambda_3, \lambda_4, \lambda_5$.

	x_0	x_1	x_2	x_3	x_4	x_5	W_i	μ_i
λ_1	0	0	0	-3	1	1	pt	-1
λ_2	0	-1	0	-1	0	1	pt	-1
λ_3	0	0	-1	-1	1	0	pt	-1
λ_4	0	-1	-1	1	0	0	pt	-1
λ_5	-1	-1	-1	0	0	0	pt	-3

TABLE 1. The weights and wall data associated to each one-parameter subgroup

Let us first justify the computation of the wall data (W_2, μ_2) in the case of the wall τ_2 determined by λ_2. The anticanonical divisor on X is given by the sum of the D_i. In the basis D_0, D_3, D_4, D_5 we can sum the columns of γ in Figure 9 to obtain

$$-K_X = -3D_0 + D_3 + 2D_4 + 2D_5.$$

Therefore we can take the dot product of $\lambda_2 = (0, -1, 0, 1)$ with the anticanonical $(3, -1, -2, -2)$ to obtain $\mu_2 = -1$. In the same manner, one can calculate each of the μ_i

Now recall that W_2 is defined as the GIT quotient of the fixed locus of λ_2 by $(\mathbb{G}_m)^3$ determined by a character α in the relative interior of τ_2. From Table 1, we see that the fixed locus of λ_2 is defined by the x_i of weight zero, namely x_0, x_2, x_4. This yields a GIT quotient of $\mathbb{C}^3$ by $(\mathbb{G}_m)^3$. Under change of basis, the action can be diagonalized completely so that weights of the action are $(1, 0, 0), (0, 1, 0)$, and $(0, 0, 1)$. Any GIT quotient with a generic character yields a point. Also in the same manner, one can see that each W_i is a point.

Let us consider the application of Theorem 3.1.5 to the wall-crossing associated to λ_2 with wall data $(pt, -1)$. For each $d \in \mathbb{Z}$, we get fully-faithful functors

$$\Phi_d : \mathrm{D}^{\mathrm{b}}(\mathrm{coh}\, Y) \to \mathrm{D}^{\mathrm{b}}(\mathrm{coh}\, X)$$

and

$$\Upsilon_+ : \mathrm{D}^{\mathrm{b}}(\mathrm{coh}\, pt) \to \mathrm{D}^{\mathrm{b}}(\mathrm{coh}\, X)$$

and a semi-orthogonal decomposition

$$\mathrm{D}^{\mathrm{b}}(\operatorname{coh} X) = \langle \mathrm{D}^{\mathrm{b}}(\operatorname{coh} pt), \mathrm{D}^{\mathrm{b}}(\operatorname{coh} Y)\rangle.$$

Fix $d = -1$. The functor Υ_+ can be described as follows. The fixed locus of λ_2 is

$$\mathbb{A}^3 = V(x_0, x_1, x_2) \subset \mathbb{A}^6.$$

The set of points,

$$\{x \in \mathbb{A}^6 \mid \lim_{t\to 0} \lambda_2(t)\cdot x \text{ exists}\} \cong \mathbb{A}^5,$$

is the vanishing locus of x_5. (Here, it useful to remember that the passing from the rings to spectra inverts the weights of an action.)

There is a sequence of morphisms of stacks,

$$\mathbb{A}^3 /\!\!/ \mathbb{G}_m^3 \to \mathbb{A}^5 /\!\!/ \mathbb{G}_m^4 \to \mathbb{A}^6 /\!\!/ \mathbb{G}_m^4 = X.$$

The first map is pulling back by the map,

$$x \mapsto \lim \lambda_2(t)\cdot x,$$

and using the projection,

$$\mathbb{G}_m^4 \cong G \to G/\lambda_2(\mathbb{G}_m) \cong \mathbb{G}_m^3.$$

The second is the inclusion.

Applying Theorem 3.1.4, we get a semi-orthogonal decomposition

$$\mathrm{D}^{\mathrm{b}}(\operatorname{coh} X) = \langle \mathcal{O}_{D_5}(-K_X), f_5^* \mathrm{D}^{\mathrm{b}}(\operatorname{coh} Y)\rangle.$$

Proceeding similarly for the walls τ_3, τ_4, τ_5 corresponding to the one-parameter subgroups $\lambda_3, \lambda_4, \lambda_5$, the exceptional collection of $\mathrm{D}^{\mathrm{b}}(\operatorname{coh} X)$ corresponding to the run of the Mori program in Figure 10 is given by

$$\langle \mathcal{O}_{D_4}(-K_X), \mathcal{O}_{D_5}(-K_X), (f_4\circ f_5)^*(\mathcal{O}_{D_3}(-3D_0 + D_3)), (g\circ f_4\circ f_5)^* \mathrm{D}^{\mathrm{b}}(\operatorname{coh}\mathbb{P}^2)\rangle.$$

Remark 5.2.1. In this way, one can also obtain a semi-orthogonal decomposition of $\mathrm{D}^{\mathrm{b}}(\operatorname{coh} X')$,

$$\langle \mathcal{O}_p(-K_{X'}), \Phi_d(\mathrm{D}^{\mathrm{b}}(\operatorname{coh} X))\rangle$$

where p is a stacky point defined by the intersection of D_4 and D_5 in X'.

It is natural for computational reasons to move the wall contributions from the left to the right. This amounts to applications of Serre functors for some of the admissible subcategories. This leaves us with

$$\mathrm{D}^{\mathrm{b}}(\operatorname{coh} X) = \langle (g\circ f_4\circ f_5)^* \mathrm{D}^{\mathrm{b}}(\operatorname{coh}\mathbb{P}^2), (f_4\circ f_5)^* \mathcal{O}_{D_3}, \mathcal{O}_{D_4}, \mathcal{O}_{D_5}\rangle.$$

Remark 5.2.2. The reader may be concerned that we mutated our exceptional before comparing it with the A-side. The difference between this choice and the original collection on the A-side amounts to a choice of orientation for the vanishing paths. It is therefore inconsequential.

Now, notice that

$$(f_4\circ f_5)^* \mathcal{O}_{D_3} = \mathcal{O}_{D_3+D_4+D_5}\,.$$

Furthermore, the unique run of the Mori program for the toric GIT description of $\mathbb{P}^2$ yields a semi-orthogonal decomposition

$$\mathrm{D}^{\mathrm{b}}(\operatorname{coh}\mathbb{P}^2) = \langle \mathcal{O}(-2), \mathcal{O}(-1), \mathcal{O}\rangle.$$

or

$$(g \circ f_4 \circ f_5)^* \, \mathrm{D}^{\mathrm{b}}(\mathrm{coh}\, \mathbb{P}^2) = \langle \mathcal{O}(-2D_0), \mathcal{O}(-D_0), \mathcal{O} \rangle.$$

In summary we have an exceptional collection $\langle E_0, \ldots, E_5 \rangle$ on $\mathrm{D}^{\mathrm{b}}(\mathrm{coh}\, X)$ where

$$\begin{aligned} E_0 &= \mathcal{O}(-2D_0), \\ E_1 &= \mathcal{O}(-D_0), \\ E_2 &= \mathcal{O}, \\ E_3 &= \mathcal{O}_{D_3+D_4+D_5}, \\ E_4 &= \mathcal{O}_{D_4}, \\ E_5 &= \mathcal{O}_{D_5}. \end{aligned}$$

Our aim is to compute the A_∞-algebra

$$A = \mathrm{End}^\bullet \left(\oplus_{i=0}^5 E_i \right) \tag{5.9}$$

As a first step, we compute the underlying cohomology algebra which gives the quiver with relations in Figure 12.



FIGURE 12. Scissors quiver

In Figure 12 the degree of each arrow is zero except

$$\deg(d_4) = 1 = \deg(d_5). \tag{5.10}$$

The relations are

$$\begin{aligned} b_2 a_1 &= b_1 a_2, \\ b_2 a_0 &= b_0 a_2, \\ b_1 a_0 &= b_0 a_1, \\ t b_1 &= 0 = t b_2, \\ d_4 t &= 0 = d_5 t. \end{aligned} \tag{5.11}$$

As a formal consequence of the homological grading on the A_∞ higher products, it is possible to eliminate all such higher products except m_3^A. To compute m_3, we must resolve the sheaves by either injective sheaves, or by the total complex of a vector bundle and Čech resolution (see also Remark 5.1.7). However, in this case we may perform one left mutation to obtain another exceptional collection

$$\langle \tilde{E}_0, \tilde{E}_1, \tilde{E}_2, \tilde{E}_3, \tilde{E}_4, \tilde{E}_5 \rangle = \langle E_0, E_1, L_{E_2} E_3, E_2, E_4, E_5 \rangle \tag{5.12}$$

where the twist can be written concretely as $L_{E_2} E_3 = \mathcal{O}(-D_1 - D_2 - D_3)$. One can check that this exceptional collection is strong so its endomorphim algebra is formal.

To get a controlled dg-resolution of A, we may replace $\mathcal{O}_{D_3+D_4+D_5}$ by the complex,

$$\mathcal{O}(-D_3 - D_4 - D_5) \xrightarrow{x_3x_4x_5} \mathcal{O},$$

and consider the resulting dg-quiver in Figure 13.



FIGURE 13. Resolved scissors quiver

All arrows are degree zero except

$$\deg(d_4) = \deg(d_5) = 1 \tag{5.13}$$

$$\deg(\tilde{b_1}) = \deg(\tilde{b_2}) = -1. \tag{5.14}$$

The relations on arrows in the quiver illustrated in Figure 13 are

$$\begin{gathered} b_j a_i = b_i a_j \ 0 \le i, j \le 2 \\ \tilde{b}_1 a_2 = \tilde{b}_2 a_1 \\ d_j \tilde{b}_i = 0 \ 1 \le i \le 2 \ , \ 4 \le j \le 5 \\ c_j s b_i = 0 \ 0 \le i \le 3 \ , \ 4 \le j \le 5 \\ d_j s b_i = 0 \ 1 \le i \le 2 \ , \ 4 \le j \le 5 \\ c_4 \tilde{b}_2 = d_4 s b_0 \ , \ c_5 \tilde{b}_1 = d_5 s b_0 \\ c_4 \tilde{b}_1 = c_5 \tilde{b}_2 = 0 \end{gathered} \tag{5.15}$$

The differential is zero on all arrows except

$$d(\tilde{b}_i) = s b_i, 1 \le i \le 2. \tag{5.16}$$

The differential then extends linearly over the path algebra.

To obtain the higher morphisms on A, one implements homological perturbation (see Section 2.1)

$$h \circlearrowright \widehat{A} \underset{i}{\overset{\pi}{\rightleftarrows}} A$$

where i is the inclusion using the already enforced notational identification and setting $i(t) = s$. The map π is the obvious projection and h is the homotopy which vanishes except for

$$h(sb_i) = \tilde{b}_i \ , \ 1 \leq i \leq 2 \tag{5.17}$$

$$h(sb_ia_j) = h(sb_ja_i) = \tilde{b}_ia_j \ , \ 1 \leq i \leq 2, 0 \leq j \leq 3. \tag{5.18}$$

To calculate m_3^A we use the formula (2.2) obtained by homological perturbation in Section 2.1 which we recall here.

$$m_3^A(x,y,z) = \pi(m_{\tilde{A}}^2(h(m_2^A(i(x),i(y))),i(z))) - (-1)^{|x|}m_{\tilde{A}}^2(i(x),h(m_2^A(i(y),i(z))))). \tag{5.19}$$

This can be computed to give zero on all triples other than three possible sequences of elements in A. We obtain

$$m_3^A(d_4,t,b_2) = c_4tb_0, \qquad m_3^A(d_5,t,b_1) = c_5tb_0, \tag{5.20}$$

$$m_3^A(d_4,t,b_2a_0) = c_4tb_0a_0, \qquad m_3^A(d_5,t,b_1a_0) = c_5tb_0a_0 \tag{5.21}$$

$$m_3^A(d_4,tb_0,a_2) = c_4tb_0a_0, \qquad m_3^A(d_5,tb_0,a_1) = c_5tb_0a_0 \tag{5.22}$$

5.2.2. *The A-model computation.* The mirror picture of the run of the toric Mori program appearing in Figure 10 follows from the monotone path polytope description. It is a degenerated LG model $\{\psi(t)\}$. The cycle corresponding to the maximal degeneration $\psi(0)$ has a sequence of components $\{Z_1, Z_2, Z_3, Z_4, Z_5\}$ which are weighted projective line orbits in $\mathcal{X}_{\Sigma(A)}$. Combinatorially, this corresponds to a path in $\Sigma(A)$ with vertices $P = \{T_0, \ldots, T_5\}$ associated to triangulations of (Q, A) and edges $\{T_i, T_{i+1}\}$ associated to extended circuits. Note that this is just a reinterpretation of the choice of a run of the toric Mori program.

Utilizing the formula given in (4.13), the ramifications of the components $\psi_i(0)$ of $\psi(0)$ can be computed using the vertices ϕ_{T_i} to be $\{3,1,1,1,1\}$. Indeed, this formula amounts to finding the difference between the sum of the volumes of all triangles containing 0 as a vertex in T_{i+1} with the same sum for T_i. Inspecting Figure 14, one sees that no triangles contain 0 in T_0, while three contain it T_1 yielding $\langle e_0, \varphi_{T_1} - \varphi_{T_0}\rangle = 3$. For each remaining edge, a single unit volume triangle is added to the set containing zero as a vertex, so each has multiplicity $a_i = 1$. As there are no degenerate circuits in this sequence, this also gives the number of critical values in each annular decomposition of $\psi(t)$ for $t \approx 0$.

Figure 14 illustrates, the triangulations T_i, together with the critical fibers arranged into annuli as t approaches zero and the tropical limit of the hypersurface over $\psi(t)$ near the bounding radii of each annulus. Each critical fiber is color-coordinated with the corresponding critical value.

As a sanity check with the B-side, let us pause for a moment to consider Figure 15 which displays the tropicalization of the fiber corresponding to the outer radius of the annulus associated to Z_4 (or equivalently the tropical variety dual to T_5). Notice that the two vanishing cycles (pictured in red and blue) are disjoint. Applying Proposition 5.1.5, one confirms immediately that the semi-orthogonal components $\mathrm{Fuk}^{\rightarrow}(\psi_3(t))$ and $\mathrm{Fuk}^{\rightarrow}(\psi_4(t))$ are in fact orthogonal. This is certainly the case on the B-side, as these two categories are generated by structure sheaves of disjoint exceptional loci.

To each critical fiber in Figure 14 we also associate a color-coordinated vanishing cycle in Figure 16. This would be done, theoretically, using the prescription in Section 5.1.1 which characterizes the vanishing cycles associated to a circuit as stable tropical Morse cells.



FIGURE 14. The potential near the mirror degeneration.



FIGURE 15. The mirror fiber and tropical vanishing cycles for X.

However, as the calculus mentioned in Section 5.1 has not yet been developed, we instead utilize the local model for vanishing cycles of surface circuits obtained in Figure 7.

Near the first component Z_0 in the degeneration, $\psi(t)$ has 3 critical values whose vanishing cycle consists of the $(1,3)$ circuit vanishing cycle L_2 drawn in Figure 7 along with two vanishing cycles. The first of these, L_1, is obtained by Dehn twisting once around each cylinder corresponding to an internal edge of the stable complex and the second, L_0, is obtained by Dehn twisting twice along these cylinders. For each of the remaining critical values, we obtain the local model of a $(2,2)$ circuit. As the vanishing cycle flows along its distinguished basis, one must keep careful track of its image in the fiber. Assembling these models onto the fiber near the outer radii of the Z_4 degeneration produces Figure 16.

We conclude with a visual inspection of the A_∞-algebra associated to the cycles $\oplus_{i=0}^5 L_i$ in $\mathrm{Fuk}^{\rightarrow}(\psi)$ as illustrated in Figure 16. To this end, we have labeled the intersections of the vanishing cycles in bijection with the morphisms obtained in the previous section for the B-model.



FIGURE 16. The fiber F with vanishing cycles and one example of a higher product

For the rigorous definition of the category associated to a collection of vanishing cycles in the general setting of symplectic Lefschetz pencils, see [Sei08]. We briefly recall the definition of the higher products m_k now .

The A_∞-structure maps $\{m_k\}_{k\geq 1}$ are determined by a count of holomorphic polygons with appropriate boundary conditions. More precisely, let $L_{i_k}, \ldots, L_{i_1}$ be k distinct vanishing cycles in the collection, with $i_1 < \cdots < i_k$. Here the subscript indicates to which path in the distinguished basis the vanishing cycle corresponds. By a $(k+1)$-marked disc Δ, we mean the disc in $\mathbb{C}$ with $(k+1)$ marked points $\{p_0, p_1, \ldots, p_k\}$ on the boundary oriented counterclockwise. We think of the k points p_i as incoming points and p_0 as an outgoing point and write $\partial_i\Delta$ for the arc on the boundary of Δ between p_i and p_{i+1}.

Finally, take $\mathcal{M}_k(F, L_{i_k}, \dots L_{i_0})$ to be the moduli space of holomorphic maps u from a $(k+1)$-marked disc Δ to F such that $u(\partial_j \Delta) \subset L_{i_j}$. Let $\mathcal{M}_k^0(F, L_{i_k}, \dots L_{i_0})$ be the zero dimensional subspace in $\mathcal{M}_k(F, L_{i_k}, \dots L_{i_0})$. In general, this space is determined by considering maps with Maslov index 2, which in the 1 dimensional setting amounts to restricting to holomorphic immersions. For exact manifolds like F, the zero dimensional moduli spaces are compact and can be given an orientation, so that we obtain an integer $\chi(\mathcal{M}_k(F, L_{i_k}, \dots L_{i_0})) \in \mathbb{Z}$ equal to the Euler characteristic. Then for $p_{i_j} \in L_{i_j} \cap L_{i_{j+1}} = \mathrm{Hom}^*(L_{i_j}, L_{i_{j+1}})$ we compute the k-th A_∞-product as

$$m_k(a_{i_k}, \dots, p_{i_0}) = \sum_{p_0 \in L_{i_k} \cap L_{i_0}} \chi(\mathcal{M}_k(F, L_{i_k}, \dots L_{i_0}))p_0. \tag{5.23}$$

Applying this formula to the vanishing cycles $\{L_0, \dots, L_5\}$ in the fiber X, one sees the underlying scissors quiver in Figure 12 based on the marking of arrows in Figure 14. To verify the relations obtained for the B-model, one must count holomorphic triangles according to (5.23). One such count is made between the Lagrangians L_0, L_1, L_2 in Figure 14. It produces the relation

$$m_2^A(b_2, a_1) = m_2^A(a_2, b_1).$$

The rest of the relations are handled similarly.

To calculate the higher products m_3 one verifies that there are six holomorphic quadralaterals with cyclic boundary conditions. One such relation and one quadralateral appears in Figure 16 between the Langrangians L_0, L_2, L_3, L_4. It produces higher product

$$m_3^A(c_4, t, b_2 a_0) = d_4 t b_0 a_0.$$

The rest can be found at www.math.miami.edu/ gdkerr/products.pdf. Finally, $m_n = 0$ for $n \geq 4$ as a formal consequence of the underlying cohomology algebra just like on the B-side.

References

[Abo09] M. Abouzaid. *Morse homology, tropical geometry, and homological mirror symmetry for toric varieties*. Selecta Math. (N.S.) 15 (2009), no. 2, 189-270.

[AFOOO] M. Abouzaid, K. Fukaya, Y.-G. Oh, H. Ohta, K. Ono. *Quantum cohomology and split generation in Lagrangian Floer theory*. In preparation.

[AS10] M. Abouzaid, P. Seidel. *An open string analogue of Viterbo functoriality*. Geom. Topol. 14 (2010), no. 2, 627-718.

[Ale02] V. Alexeev. *Complete moduli in the presence of semiabelian group action*. Ann. of Math. (2) 155 (2002), no. 3, 611-708

[AGM94] P. Aspinwall, B. Greene, D. Morrison. *Calabi-Yau Moduli Space, Mirror Manifolds and Spacetime Topology Change in String Theory*. Nucl.Phys. B416 (1994) 414-480.

[AKO06] D. Auroux, L. Katzarkov, D. Orlov. *Mirror symmetry for del Pezzo surfaces: vanishing cycles and coherent sheaves*. Invent. Math. 166 (2006), no. 3, 537-582.

[AKO08] D. Auroux, L. Katzarkov, D. Orlov. *Mirror symmetry for weighted projective planes and their noncommutative deformations*. Ann. of Math. (2) 167 (2008), no. 3, 867-943.

[BFK12] M. Ballard, D. Favero, L. Katzarkov. *Variation of Geometric Invariant Theory quotients and derived categories*. arXiv:1203.6643v2.

[Bei78] A. Beilinson. *Coherent sheaves on $\mathbb{P}^n$ and problems in linear algebra*. Functional Anal. Appl. 12 (1978), no.3, 214-216.

[BS92] L. Billera, B. Sturmfels. *Fiber Polytopes*. Ann. of Math. 135 (1992), no. 3, 527-549.

[Bon89] A. Bondal. *Representations of associative algebras and coherent sheaves*. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 53 (1989), no. 1, 25-44; translation in Math. USSR-Izv. 34 (1990), no. 1, 23-42.

[BK89] A. Bondal, M. Kapranov. *Representable functors, Serre functors, and reconstructions.* (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 53 (1989), no. 6, 1183-1205, 1337; translation in Math. USSR-Izv. 35 (1990), no. 3, 519-541.

[BR] A. Bondal, W.-D. Ruan. *Mirror symmetry for weighted projective spaces.* In preparation.

[BV03] A. Bondal, M. Van den Bergh. *Generators and representability of functors in commutative and non-commutative geometry.* Mosc. Math. J. 3 (2003), no.1, 1-36.

[BCS05] L. Borisov, L. Chen, G. Smith. *The orbifold Chow ring of toric Deligne-Mumford stacks.* J. Amer. Math. Soc. 18 (2005), no. 1, 193-215.

[CL12] K. Chan, N. C. Leung. *Matrix factorizations from SYZ transformations.* in Advances in Geometric Analysis, 203-224, Adv. Lect. Math. (ALM) 21, Int. Press, Somerville, MA, 2012.

[CS12] Q. Chen, M. Satriano. *Chow quotients of toric varieties as moduli space of log stable maps.* arXiv:1201.3406.

[CO06] C.-H. Cho, Y.-G. Oh, *Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds*, Asian J. Math. 10 (2006), 773-814.

[Cox98] D. Cox. *Recent developments in toric geometry.* Algebraic Geometry - Santa Cruz 1995. Proc. Sympos. Pure Math., 62 Part 2. American Mathematical Society. 1998.

[CLS11] D. Cox, J. Little, H. Schenck. Toric varieties. Graduate Studies in Mathematics, 124. American Mathematical Society, Providence, RI, 2011.

[DKK12] C. Diemer, L. Katzarkov, G. Kerr. *Symplectomorphism group relations and degenerations of Landau-Ginzburg models.* arXiv:1204.2233.

[DH98] I. Dolgachev, Y. Hu. *Variation of Geometric Invariant Theory quotients.* With an appendix by N. Ressayre. Inst. Hautes Études Sci. Publ. Math. No. 87 (1998), 5-56.

[FHKV09] B. Feng, Y.-H. He, K. Kennaway, C. Vafa. *Dimer models from mirror symmetry and quivering amoebae.* Adv. Theor. Math. Phys. 12 (2008), no. 3, 489-545.

[FLTZ08] B. Fang, C.-C. M. Liu, D. Treumann, E. Zaslow. *T-duality and Homological Mirror Symmetry for toric varieties.* Adv. Math. 229 (2012), no. 3, 1875-1911.

[GKZ94] I. Gel'fand, M. Kapranov, A. Zelevinsky. Discriminants, resultants and multidimensional determinants. Reprint of the 1994 edition. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, 2008.

[GS11] A. Geraschenko, M. Satriano. *Toric stacks I: The theory of stacky fans.* arXiv:1107.1906.

[GKH11] M. Gross, S. Keel, P. Hacking. *Mirror symmetry for log Calabi-Yau surfaces I.* arXiv:1106.4977.

[Hac03] P. Hacking. *Compact moduli of hyperplane arrangements.* arXiv:math/0310479.

[H-L12] D. Halpern-Leistner. *The derived category of a GIT quotient.* arXiv:1203.0276.

[HV02] K. Hori, C. Vafa. *Mirror symmetry.* arXiv:hep-th/0002222.

[HK99] Y. Hu, S. Keel. *A GIT proof of Włodarczyk's weighted factorization theorem.* arXiv:math.AG/9904146.

[HK00] Y. Hu, S. Keel. *Mori dream spaces and GIT.* Dedicated to William Fulton on the occasion of his 60th birthday. Michigan Math. J. 48 (2000), 331-348.

[Kap95] M. Kapranov. *Chow quotients of Grassmannians I.* I.M. Gel'fand seminar, 29-110. Adv. Soviet Math., 16, Part 2. Amer. Math Soc. 1993.

[KSZ91] M. Kapranov, A. Zelevinsky, B. Sturmfels. *Quotients of toric varieties.* Math. Ann. 290 (1991), no. 4, 643-655.

[KKP13] L. Katzarkov, M. Kontsevich, T. Pantev. *On complex moduli for Landau-Ginzburg models.* In preparation.

[Kaw02] Y. Kawamata. *D-equivalence and K-equivalence.* J. Differential Geom. 61 (2002), no. 1, 147-171.

[Kaw05] Y. Kawamata. *Log crepant birational maps and derived categories.* J. Math. Sci. Univ. Tokyo 12 (2005), no. 2, 211-231.

[Kaw06] Y. Kawamata. *Derived categories of toric varieties.* Michigan Math. J. 54 (2006), no. 3, 517-535.

[Kaw12] Y. Kawamata. *Derived categories of toric varieties, II.* arXiv:1201.3460.

[Kee92] S. Keel. *Intersection theory of moduli space of stable n-pointed curves of genus zero.* Trans. Amer. Math. Soc. 330 (1992), no. 2, 545-574.

[Kel94] B. Keller. *Deriving DG categories.* Ann. Sci. cole Norm. Sup. (4) 27 (1994), no. 1, 63-102.

[Kel01] B. Keller. *Introduction to A-infinity algebras and modules.* Homology Homotopy Appl. 3 (2001), no. 1, 1-35.

[Kon95] M. Kontsevich. *Homological algebra of mirror symmetry.* Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zrich, 1994), 120-139, Birkhuser, Basel, 1995.
[Kon98] M. Kontsevich. Course at ENS, 1998.
[Laf03] L. Lafforgue. *Chirurgie des grassmanniennes.* CPM Monograph Series. vol 19. American Mathematical Society. 2003.
[L-H03] K. Lefèvre-Hasegawa. *Sur le A-infini catégories.* arXiv:math/0310337.
[Mar04] M. Markl, Martin. *Transferring A_∞ (strongly homotopy associative) structures.* Rend. Circ. Mat. Palermo (2) Suppl. No. 79 (2006), 139-151.
[Mat02] K. Matsuki. *Introduction to the Mori Program.* Universitext. Springer-Verlag, New York, 2002.
[Mik04] G. Mikhalkin. *Decomposition into pairs-of-pants for complex algebraic hypersurfaces.* Topology. Vol 43, p. 1035-1065. 2004.
[MiSt05] E. Miller, B. Sturmfels. Combinatorial commutative algebra. Graduate Texts in Mathematics, 227. Springer-Verlag, New York, 2005.
[MFK94] D. Mumford, J. Fogarty, F. Kirwan. Geometric Invariant Theory. Third edition. Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)], 34. Springer-Verlag, Berlin, 1994.
[OP91] T. Oda, H.S. Park. *Linear Gale transforms and Gelfand-Kapranov-Zelevinskij decompositions.* Tohoku Math. J. (2) 43 (1991), no. 3, 375-399.
[Rei83] M. Reid. *Decomposition of toric morphisms.* Arithmetic and geometry, Vol. II, 395-418, Progr. Math., 36, Birkhäuser Boston, Boston, MA, 1983.
[Res00] N. Ressayre. *The GIT-equivalence for G-line bundles.* Geom. Dedicata 81 (2000), no. 1-3, 295-324.
[Sei00] P. Seidel. *More about vanishing cycles and mutation.* Symplectic geometry and mirror symmetry (Seoul, 2000), 429-465, World Sci. Publ., River Edge, NJ, 2001.
[Sei08] P. Seidel. *Fukaya categories and Picard-Lefschetz theory.* Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.
[Sei11] P. Seidel. *Some speculations on pair-of-pants decompositions and Fukaya categories.* arXiv:1004.0906.
[She11] N. Sheridan. *On the homological mirror symmetry conjecture for pairs of pants.* J. Differential Geom. 89 (2011), no. 2, 271-367.
[Sta63] J. Stasheff. *Homotopy associativity of H-spaces. I, II.* Trans. Amer. Math. Soc. 108 (1963), 275-292; ibid. 108 (1963) 293-312.
[SYZ96] A. Strominger, S.-T. Yau, E. Zaslow *Mirror symmetry is T-duality.* Nuclear Phys. B 479 (1996), no. 1-2, 243-259.
[Tha96] M. Thaddeus. *Geometric Invariant Theory and flips.* J. Amer. Math. Soc. 9 (1996), no. 3, 691-723.
[Ued06] K. Ueda. *Homological mirror symmetry for toric del Pezzo surfaces.* Comm. Math. Phys. 264 (2006), no. 1, 71-85.
[UY07] K. Ueda, M. Yamazaki. *Homological mirror symmetry for toric orbifolds of toric del Pezzo surfaces.* J. reine angew. Math. Ahead of Print: DOI 10.1515/crelle.2012.031.

Matthew Ballard
University of Wisconsin-Madison, Department of Mathematics, Madison, WI, USA
Universität Wien, Fakultät für Mathematik, Wien, Österreich
Email: **ballard@math.wisc.edu**

Colin Diemer
University of Miami, Department of Mathematics, Coral Gables, FL, USA
Universität Wien, Fakultät für Mathematik, Wien, Österreich
Email: **cdiemer@gmail.com**

David Favero
Universität Wien, Fakultät für Mathematik, Wien, Österreich
Email: **favero@gmail.com**

Ludmil Katzarkov
University of Miami, Department of Mathematics, Coral Gables, FL, USA
Universität Wien, Fakultät für Mathematik, Wien, Österreich
Email: **lkatzark@math.uci.edu**

Gabriel Kerr
University of Miami, Department of Mathematics, Coral Gables, FL, USA
Email: **gabriel.d.kerr@gmail.com**